

PRIME-MD & PHQ Family of Scales Reference List by Author (Updated April 2010)

- (1) Self-reported illness and health status among Gulf War veterans. A population-based study. The Iowa Persian Gulf Study Group. *JAMA*. 1997;277:238-245.
- (2) Research Validates Tool for Assessing Mental Disorders among Adolescent Primary Care Patients. MCH Alert. National Center for Education in Maternal and Child Health . 2002.
Ref Type: Electronic Citation
- (3) Patient Health Questionnaire for Adolescents (PHQ-A) Release Announcement. Center for Promotion of Child Development through Primary Care CHADIS website. 2007.
Ref Type: Electronic Citation
- (4) Abdel-Kader K, Unruh ML, Weisbord SD. Symptom Burden, Depression, and Quality of Life in Chronic and End-Stage Kidney Disease. *Clinical Journal of the American Society of Nephrology*. 2009;4:1057-1064.
- (5) Ackermann RT, Rosenman MB, Downs SM et al. Telephonic case-finding of major depression in a Medicaid chronic disease management program for diabetes and heart failure. *General Hospital Psychiatry*. 2005;27:338-343.
- (6) Acton GS, Prochaska JJ, Kaplan AS, Small T, Hall SM. Depression and stages of change for smoking in psychiatric outpatients. *Addict Behav*. 2001;26:621-631.
- (7) Adewuya AO, Ola BA, Afolabi OO. Validity of the patient health questionnaire (PHQ-9) as a screening tool for depression amongst Nigerian university students. *J Affect Disord*. 2006;96:89-93.
- (8) Adler DA, McLaughlin TJ, Rogers WH, Hong C, Lapitsky L, Lerner D. Job performance deficits due to depression. *American Journal of Psychiatry*. 2006;163:1569-1576.
- (9) Aguera L, Failde I, Cervilla JA, az-Fernandez P, Mico JA. Medically unexplained pain complaints are associated with underlying unrecognized mood disorders in primary care. *Bmc Family Practice*. 2010;11.
- (10) Aiarzaguena JM, Grandes G, Salazar A, Gaminde I, Sanchez A. The diagnostic challenges presented by patients with medically unexplained symptoms in general practice. *Scandinavian Journal of Primary Health Care*. 2008;26:99-105.
- (11) Al-Windi A. The relations between symptoms, somatic and psychiatric conditions, life satisfaction and perceived health. A primary care based study. *Health Qual Life Outcomes*. 2005;3:28.
- (12) Al-Windi A. Depression in general practice. *Nordic Journal of Psychiatry*. 2005;59:272-277.
- (13) Albert NM, Moser DK, Nutter B, Pozuelo L. Are PHQ-9 and PHQ-2 Depression Score Cutoffs the Best Cutoffs for Determining Significant Depression in Pts with HF and Mild-Moderate Symptoms? *Journal of Cardiac Failure*. 2009;15:384.

- (14) Alvanzo AAH, Svikis DS. History of physical abuse and periconceptional drinking in pregnant women. *Substance Use & Misuse*. 2008;43:1098-1109.
- (15) Amin AA, Menon RA, Reid KJ, Harris WS, Spertus JA. Acute Coronary Syndrome Patients With Depression Have Low Blood Cell Membrane Omega-3 Fatty Acid Levels. *Psychosomatic Medicine*. 2008;70:856-862.
- (16) Anastario MP, Larrance R, Lawry L. Using Mental Health Indicators to Identify Postdisaster Gender-Based Violence among Women Displaced by Hurricane Katrina. *Journal of Womens Health*. 2008;17:1437-1444.
- (17) Andersson L, Sundstrom-Poromaa I, Bixo M, Wulff M, Bondestam K, Astrom M. Point prevalence of psychiatric disorders during the second trimester of pregnancy: A population-based study. *American Journal of Obstetrics and Gynecology*. 2003;189:148-154.
- (18) Andersson L, Sundstrom-Poromaa I, Wulff M, Astrom M, Bixo M. Neonatal outcome following maternal antenatal depression and anxiety: A population-based study. *American Journal of Epidemiology*. 2004;159:872-881.
- (19) Andersson L, Sundstrom-Poromaa I, Wulff M, Astrom M, Bixo M. Implications of antenatal depression and anxiety for obstetric outcome. *Obstetrics and Gynecology*. 2004;104:467-476.
- (20) Ani C, Bazargan M, Hindman D et al. Depression symptomatology and diagnosis: discordance between patients and physicians in primary care settings. *Bmc Family Practice*. 2008;9.
- (21) Annunziato RA, Rubinstein D, Murgueitio M et al. Psychiatric Symptom Presentation in Ethnically Diverse Cardiology Patients. *Ethnicity & Disease*. 2009;19:271-275.
- (22) Ansseau M, Dierick M, Buntinx F et al. High prevalence of mental disorders in primary care. *Journal of Affective Disorders*. 2004;78:49-55.
- (23) Arean PA, Ayalon L. Assessment and treatment of depressed older adults in primary care. *Clinical Psychology-Science and Practice*. 2005;12:321-335.
- (24) Arnau RC, Meagher MW, Norris MP, Bramson R. Psychometric evaluation of the Beck Depression Inventory-II with primary care medical patients. *Health Psychol*. 2001;20:112-119.
- (25) Arnow BA, Hunkeler EM, Blasey CM et al. Comorbid depression, chronic pain, and disability in primary care. *Psychosomatic Medicine*. 2006;68:262-268.
- (26) Arnow BA, Blasey CM, Lee J et al. Relationships Among Depression, Chronic Pain, Chronic Disabling Pain, and Medical Costs. *Psychiatric Services*. 2009;60:344-350.
- (27) Avasthi A, Varma SC, Kulhara P, Nehra R, Grover S, Sharma S. Diagnosis of common mental disorders by using PRIME-MD Patient Health Questionnaire. *Indian Journal of Medical Research*. 2008;127:159-164.
- (28) Avasthi A, Varma SC, Kulhara P, Nehra R, Grover S, Sharma S. PRIME-MD health questionnaire for diagnosis of common mental disorders - Response. *Indian Journal of Medical Research*. 2008;127:506.

- (29) Baca E, Saiz J, Aguera L et al. Validation of the Spanish version of PRIME-MD: a procedure for diagnosing mental disorders in primary care. *Actas Esp Psiquiatr*. 1999;27:375-383.
- (30) Backenstrass M, Frank A, Joest K, Hingmann S, Mundt C, Kronmuller KT. A comparative study, of nonspecific depressive symptoms and minor depression regarding functional impairment and associated characteristics in primary care. *Comprehensive Psychiatry*. 2006;47:35-41.
- (31) Bailer J, Schwarz D, Witthoft M, Stubinger C, Rist F. Prevalence of Mental Disorders Among College Students at a German University. *Psychotherapie Psychosomatik Medizinische Psychologie*. 2008;58:423-429.
- (32) Bailer J, Stubinger C, Dressing H, Gass P, Rist F, Kuhner C. Increased Prevalence of Problematic Alcohol Consumption in University Students. *Psychotherapie Psychosomatik Medizinische Psychologie*. 2009;59:376-379.
- (33) Bakas T, Kroenke K, Plue LD, Perkins SM, Williams LS. Outcomes among family caregivers of aphasic versus nonaphasic stroke survivors. *Rehabilitation Nursing*. 2006;31:33-42.
- (34) Bakas T, Champion V, Perkins SM, Farran CJ, Williams LS. Psychometric testing of the revised 15-item Bakas Caregiving Outcomes Scale. *Nursing Research*. 2006;55:346-355.
- (35) Bakker IM, Terluin B, van Marwijk HWJ, van Mechelen W, Stalman WAB. Test-retest reliability of the PRIME-MD: limitations in diagnosing mental disorders in primary care. *European Journal of Public Health*. 2009;19:303-307.
- (36) Baldomero EB, Ruiz JS, Chavarino AP. The detection of mental disorders by physicians who are not psychiatrists: Usefulness of the PRIME-MD questionnaire. *Medicine Clinica*. 2001;116:504-509.
- (37) Baldomero EB, Arrate MLC, Perez-Rodriguez MM, Baca-Garcia E. Mental disorders in victims of terrorism and their families. *Medicina Clinica*. 2004;122:681-685.
- (38) Balestrieri M, Carta MG, Leonetti S, Sebastiani G, Starace F, Bellantuono C. Recognition of depression and appropriateness of antidepressant treatment in Italian primary care. *Social Psychiatry and Psychiatric Epidemiology*. 2004;39:171-176.
- (39) Balestrieri M, Marcon G, Samani F et al. Mental disorders associated with benzodiazepine use among older primary care attenders - A regional survey. *Social Psychiatry and Psychiatric Epidemiology*. 2005;40:308-315.
- (40) Bambauer KZ, Locke SE, Aupont O, Mullan MG, McLaughlin TJ. Using the Hospital Anxiety and Depression Scale to screen for depression in cardiac patients. *General Hospital Psychiatry*. 2005;27:275-284.
- (41) Bandelow B, Brassler M. Clinical suitability of GAD-7 scale compared to hospital anxiety and depression scale-A for monitoring treatment effects in generalized anxiety disorder. *European Neuropsychopharmacology*. 2009;19:S604-S605.

- (42) bdel-Kader K, Unruh ML, Weisbord SD. Symptom Burden, Depression, and Quality of Life in Chronic and End-Stage Kidney Disease. *Clinical Journal of the American Society of Nephrology*. 2009;4:1057-1064.
- (43) Beck AT, Steer RA, Ball R, Ciervo CA, Kabat M. Use of the Beck Anxiety and Depression Inventories for Primary Care with medical outpatients. *Assessment*. 1997;4:211-219.
- (44) Beck AT, Guth D, Steer RA, Ball R. Screening for major depression disorders in medical inpatients with the Beck Depression Inventory for Primary Care. *Behav Res Ther*. 1997;35:785-791.
- (45) Becker S, Al Zaid K, Al Faris E. Screening for somatization and depression in Saudi Arabia: A validation study of the PHQ in primary care. *International Journal of Psychiatry in Medicine*. 2002;32:271-283.
- (46) Becker SM. Detection of somatization and depression in primary care in Saudi Arabia. *Social Psychiatry and Psychiatric Epidemiology*. 2004;39:962-966.
- (47) Benedikt G, Schulmeister E, Sattel H et al. Somatic symptoms and illness worries in primary care - A study of 1751 general practice attenders. *Zeitschrift fur Psychiatrie Psychologie und Psychotherapie*. 2007;55:21-29.
- (48) Bennett IM, Coco A, Coyne JC et al. Efficiency of a two-item pre-screen to reduce the burden of depression screening in pregnancy and postpartum: An IMPLICIT network study. *Journal of the American Board of Family Medicine*. 2008;21:317-325.
- (49) Benninghoven D, Specht T, Kunzendorf S et al. The Luebeck Interview for Psychosocial Screening (LIPS) - A validation study with patients suffering from coronary artery disease. *Psychotherapie Psychosomatik Medizinische Psychologie*. 2003;53:267-274.
- (50) Bensenor IM, Pereira AC, Tannuri AC et al. Systemic arterial hypertension and psychiatric morbidity in the outpatient care setting of a tertiary hospital. *Arq Neuropsiquiatr*. 1998;56:406-411.
- (51) Bentley SM, Melville JL, Berry BD, Katon WJ. Implementing a clinical and research registry in obstetrics: overcoming the barriers. *General Hospital Psychiatry*. 2007;29:192-198.
- (52) Bergus GR, Hartz AJ, Noyes R et al. The limited effect of screening for depressive symptoms with the PHQ-9 in rural family practices. *Journal of Rural Health*. 2005;21:303-309.
- (53) Bermejo I, Niebling W, Mathias B, Harter M. Patients' and physicians' evaluation of the PHQ-D for depression screening. *Primary Care & Community Psychiatry*. 2005;10:125-131.
- (54) Beutel ME, Glaesmer H, Wiltink J, Marian H, Brahler E. Life satisfaction, anxiety, depression and resilience across the life span of men. *Aging Male*. 2010;13:32-39.
- (55) Bijl D, van Marwijk HWJ, Beekman ATF, de Haan M, van Tilburg W. A randomized controlled trial to improve the recognition, diagnosis and treatment of major depression in elderly people in general practice: design, first results and feasibility of the West Friesland Study. *Primary Care Psychiatry*. 2003;8:135-140.

- (56) Birket-Smith M, Rasmussen A. Screening for mental disorders in cardiology outpatients. *Nordic Journal of Psychiatry*. 2008;62:147-150.
- (57) Bixo M, Sundstrom-Poromaa I, Bjorn I, Astrom M. Patients with psychiatric disorders in gynecologic practice. *Amer J Obstet Gynecol*. 2001;185:396-402.
- (58) Blalock JA, Fouladi RT, Wetter DW, Cinciripini PA. Depression in pregnant women seeking smoking cessation treatment. *Addictive Behaviors*. 2005;30:1195-1208.
- (59) Blalock JA, Robinson JD, Wetter DW, Cinciripini PM. Relationship of DSM-IV-based depressive disorders to smoking cessation and smoking reduction in pregnant smokers. *American Journal on Addictions*. 2006;15:268-277.
- (60) Bombardier CH, Richards JS, Krause JS, Tulsy D, Tate DG. Symptoms of major depression in people with spinal cord injury: Implications for screening. *Archives of Physical Medicine and Rehabilitation*. 2004;85:1749-1756.
- (61) Bombardier CH, Fann JR, Temkin N et al. Posttraumatic stress disorder symptoms during the first six months after traumatic brain injury. *Journal of Neuropsychiatry and Clinical Neurosciences*. 2006;18:501-508.
- (62) Bornstein RF, Porcerelli JH, Huprich SK, Markova T. Construct Validity of the Relationship Profile Test: Correlates of Overdependence, Detachment, and Healthy Dependency in Low Income Urban Women Seeking Medical Services. *Journal of Personality Assessment*. 2009;91:537-544.
- (63) Bosmans J, de Bruijne M, van Hout H et al. Cost-effectiveness of a disease management program for major depression in elderly primary care patients. *Journal of General Internal Medicine*. 2006;21:1020-1026.
- (64) Bost RH, Wen FK, Basso MR, Cates GR. Online tools for evaluating patient change: Statistical foundations, clinical applications, research relevance. *Rehabilitation Psychology*. 2008;53:313-320.
- (65) Bosworth HB, Parsey KS, Butterfield MI et al. Racial variation in wanting and obtaining mental health services among women veterans in a primary care clinic. *J Natl Med Assn*. 2000;92:231-236.
- (66) Bosworth HB, Butterfield MI, Stechuchak KM, Bastian LA. The relationship between self-rated health and health care service use among women veterans in a primary care clinic. *Womens Health Issues*. 2000;10:278-285.
- (67) Boulware LE, Ratner LE, Troll MU et al. Attitudes, psychology, and risk taking of potential live kidney donors: Strangers, relatives, and the general public. *American Journal of Transplantation*. 2005;5:1671-1680.
- (68) Bradley JR, Farnsworth DL. Testing for Mutual Exclusivity. *Journal of Applied Statistics*. 2009;36:1307-1314.

- (69) Brody DS, Hahn SR, Spitzer RL et al. Identifying patients with depression in the primary care setting - A more efficient method. *Arch Intern Med.* 1998;158:2469-2475.
- (70) Brody DS, Dietrich AJ, deGruy F, III, Kroenke K. The depression in primary care tool kit. *Int J Psychiatry Med.* 2000;30:99-110.
- (71) Broers T, Hodgetts G, Batic-Mujanovic O, Petrovic V, Hasanagic M, Godwin M. Prevalence of mental and social disorders in adults attending primary care centers in Bosnia and Herzegovina. *Croatian Medical Journal.* 2006;47:478-484.
- (72) Brucki SMD, Nitrini R. Subjective memory impairment in a rural population with low education in the Amazon rainforest: an exploratory study. *International Psychogeriatrics.* 2009;21:164-171.
- (73) Bundgaard S. "Prime-MD"--a glittering tool. *Ugeskr Laeger.* 1998;160:6987-6988.
- (74) Burgess A, Kunik ME, Stanley MA. Chronic obstructive pulmonary disease - Assessing and treating psychological issues in patients with COPD. *Geriatrics.* 2005;60:18-21.
- (75) Cabassa LJ, Lagomasino IT, Dwight-Johnson M, Hansen MC, Xie B. Measuring Latinos' Perceptions of Depression: A Confirmatory Factor Analysis of the Illness Perception Questionnaire. *Cultural Diversity & Ethnic Minority Psychology.* 2008;14:377-384.
- (76) Cadzow RB, Servoss TJ, Fox CH. The health status of patients of a student-run free medical clinic in inner-city Buffalo, NY. *Journal of the American Board of Family Medicine.* 2007;20:572-580.
- (77) Cadzow RB, Servoss TJ. The Association Between Perceived Social Support and Health Among Patients at a Free Urban Clinic. *Journal of the National Medical Association.* 2009;101:243-250.
- (78) Callahan CM, Boustani MA, Unverzagt FW et al. Effectiveness of collaborative care for older adults with Alzheimer disease in primary care - A randomized controlled trial. *Jama-Journal of the American Medical Association.* 2006;295:2148-2157.
- (79) Cameron IM, Crawford JR, Lawton K, Reid IC. Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care. *British Journal of General Practice.* 2008;58:32-36.
- (80) Cameron IM, Crawford JR, Lawton K et al. Assessing the validity of the PHQ-9, HADS, BDI-II and QIDS-SR16 in measuring severity of depression in a UK sample of primary care patients with a diagnosis of depression: study protocol. *Primary Care & Community Psychiatry.* 2008;13:67-71.
- (81) Cannon DS, Tiffany ST, Coon H, Scholand MB, McMahon WM, Leppert MF. The PHQ-9 as a brief assessment of lifetime major depression. *Psychological Assessment.* 2007;19:247-251.
- (82) Carballeira Y, Dumont P, Borgacci S et al. Criterion validity of the French version of Patient Health Questionnaire (PHQ) in a hospital department of internal medicine. *Psychology and Psychotherapy-Theory Research and Practice.* 2007;80:69-77.

- (83) Carrera M, Herran A, yuso-Mateos JL et al. Quality of life in early phases of panic disorder: Predictive factors. *Journal of Affective Disorders*. 2006;94:127-134.
- (84) Carson AJ, Ringbauer B, MacKenzie L, Warlow C, Sharpe M. Neurological disease, emotional disorder, and disability: they are related: a study of 300 consecutive new referrals to a neurology outpatient department. *J Neurol Neurosurg Psychiatr*. 2000;68:202-206.
- (85) Carson AJ, Ringbauer B, Stone J, McKenzie L, Warlow C, Sharpe M. Do medically unexplained symptoms matter? A prospective cohort study of 300 new referrals to neurology outpatient clinics. *J Neurol Neurosurg Psychiatr*. 2000;68:207-210.
- (86) Carson AJ, Postma K, Stone J, Warlow C, Sharpe M. The outcome of depressive disorders in neurology patients: a prospective cohort study. *Journal of Neurology Neurosurgery and Psychiatry*. 2003;74:893-896.
- (87) Carson AJ, Best S, Postma K, Stone J, Warlow C, Sharpe M. The outcome of neurology outpatients with medically unexplained symptoms: a prospective cohort study. *Journal of Neurology Neurosurgery and Psychiatry*. 2003;74:897-900.
- (88) Caska CM, Hendrickson BE, Wong MH, Ali S, Neylan T, Whooley MA. Anger expression and sleep quality in patients with coronary heart disease: findings from the Heart and Soul Study. *Psychosom Med*. 2009;71:280-285.
- (89) Cass AR, Volk RJ, Nease DE. Health-related quality of life in primary care patients with recognized and unrecognized mood and anxiety disorders. *Int J Psychiat Med*. 1999;29:293-309.
- (90) Chao DVK, Fabb WE, Pang AHT, Chan CSY, Leung T. Detection of mental disorder amongst primary care attenders using the Chinese version of the PRIME-MD (cPRIME-MD). *MediMedia Asia*. 1997.
- (91) Chen TM, Huang FY, Chang C, Chung H. Using the PHQ-9 for depression screening and treatment monitoring for Chinese Americans in primary care. *Psychiatric Services*. 2006;57:976-981.
- (92) Christensen O, Bundgaard S, Bech P. Prevalence of clinical (major) depression in general practice using the DSM-IV version of PRIME-MD. *Int J Psychiatr Clin Pract*. 2001;5:49-54.
- (93) Chwastiak LA, Gibbons LE, Ehde DM et al. Fatigue and psychiatric illness in a large community sample of persons with multiple sclerosis. *Journal of Psychosomatic Research*. 2005;59:291-298.
- (94) Clarke DM, Piterman L, Byrne CJ, Austin DW. Somatic symptoms, hypochondriasis and psychological distress: a study of somatisation in Australian general practice. *Medical Journal of Australia*. 2008;189:560-564.
- (95) Colenda CC, Legault C, Rapp SR et al. Psychiatric Disorders and Cognitive Dysfunction Among Older, Postmenopausal Women: Results From the Women's Health Initiative Memory Study. *American Journal of Geriatric Psychiatry*. 2010;18:177-185.

- (96) Collinsworth EM, Ross SR. Handbook of assessment and treatment planning for psychological disorders. *Cognitive and Behavioral Practice*. 2003;10:264-265.
- (97) Connelly JE, Wofford AB, Philbrick JT. Healthy patients who perceive poor health: Why are they worried sick? *Amer J Med Sci*. 2000;320:36-42.
- (98) Corapcioglu A, Ozer GU. Adaptation of revised Brief PHQ (Brief-PHQ-r) for diagnosis of depression, panic disorder and somatoform disorder in primary healthcare settings. *International Journal of Psychiatry in Clinical Practice*. 2004;8:11-18.
- (99) Corson K, Gerrity MS, Dobscha SK. Screening for depression and suicidality in a VA primary care setting: 2 items are better than 1 item. *American Journal of Managed Care*. 2004;10:839-845.
- (100) Coyne JC, Thombs BJ, Mitchell AJ. PHQ-9 and PHQ-2 in Western Kenya. *Journal of General Internal Medicine*. 2009;24:890.
- (101) Cutler CB, Legano LA, Dreyer BP et al. Screening for maternal depression in a low education population using a two item questionnaire. *Archives of Womens Mental Health*. 2007;10:277-283.
- (102) Dahl AA, Kruger MB, Dahl NH, Karlsson H, Von Knorring L, Stordal E. SPIFA-A presentation of the Structured Psychiatric Interview for General Practice. *Nordic Journal of Psychiatry*. 2009;63:443-453.
- (103) Daig I, Herschbach P, Lehmann A, Knoll N, Decker O. Gender and age differences in domain-specific life satisfaction and the impact of depressive and anxiety symptoms: a general population survey from Germany. *Quality of Life Research*. 2009;18:669-678.
- (104) Daly EJ, Trivedi MH, Raskin P, Grannemann BD. Screening for depression in a diabetic outpatient population. *International Journal of Psychiatry in Clinical Practice*. 2007;11:268-272.
- (105) Damush TM, Wu JW, Bair MJ, Sutherland JM, Kroenke K. Self-management practices among primary care patients with musculoskeletal pain and depression. *Journal of Behavioral Medicine*. 2008;31:301-307.
- (106) Das AK, Olfson M, Gameroff MJ et al. Screening for bipolar disorder in a primary care practice. *Jama-Journal of the American Medical Association*. 2005;293:956-963.
- (107) Davidson KW, Kupfer DJ, Bigger JT et al. Assessment and treatment of depression in patients with cardiovascular disease: National Heart, Lung, and Blood Institute working group report. *Psychosomatic Medicine*. 2006;68:645-650.
- (108) Davis GE, Yueh B, Walker E, Katon W, Koepsell TD, Weymuller EA. Psychiatric distress amplifies symptoms after surgery for chronic rhinosinusitis. *Otolaryngology-Head and Neck Surgery*. 2005;132:189-196.
- (109) Davis TM, Bush KR, Kivlahan DR, Dobie DJ, Bradley KA. Screening for substance abuse and psychiatric disorders among women patients in a VA health care system. *Psychiatric Services*. 2003;54:214-218.

- (110) Dbouk N, Arguedas MR, Sheikh A. Assessment of the PHQ-9 as a screening tool for depression in patients with chronic hepatitis C. *Digestive Diseases and Sciences*. 2008;53:1100-1106.
- (111) de Jonge P., Mangano D, Whooley MA. Differential association of cognitive and somatic depressive symptoms with heart rate variability in patients with stable coronary heart disease: findings from the Heart and Soul Study. *Psychosom Med*. 2007;69:735-739.
- (112) De Ridder D, Ost D, Bruyninckx F. The presence of Fowler's syndrome predicts successful long-term outcome of sacral nerve stimulation in women with urinary retention. *European Urology*. 2007;51:229-234.
- (113) DeJesus RS, Vickers KS, Melin GJ, Williams MD. A system-based approach to depression management in primary care using the patient health questionnaire-9. *Mayo Clinic Proceedings*. 2007;82:1395-1402.
- (114) Dezee KJ, Jackson JL, Hatzigeorgiou C, Kristo D. The Epworth sleepiness scale: relationship to sleep and mental disorders in a sleep clinic. *Sleep Medicine*. 2006;7:327-332.
- (115) Diefenbach GJ, Tolin DF, Meunier SA, Gilliam CM. Assessment of Anxiety in Older Home Care Recipients. *Gerontologist*. 2009;49:141-153.
- (116) Dietrich AJ, Oxman TE, Burns MR, Winchell CW, Chin T. Application of a depression management office system in community practice: A demonstration. *Journal of the American Board of Family Practice*. 2003;16:107-114.
- (117) Diez-Quevedo C, Rangil T, Sanchez-Planell L, Kroenke K, Spitzer RL. Validation and utility of the patient health questionnaire in diagnosing mental disorders in 1003 general hospital Spanish inpatients. *Psychosom Med*. 2001;63:679-686.
- (118) Dillard DA, Christopher D. The Southcentral foundation depression collaborative. *International Journal of Circumpolar Health*. 2007;45-53.
- (119) Dobscha SK, Gerrity MS, Corson K, Bahr A, Cuilwik NM. Measuring adherence to depression treatment guidelines in a VA primary care clinic. *General Hospital Psychiatry*. 2003;25:230-237.
- (120) Dobscha SK, Corson K, Solodky J, Gerrity MS. Use of videoconferencing for depression research: Enrollment, retention, and patient satisfaction. *Telemedicine Journal and E-Health*. 2005;11:84-89.
- (121) Dobscha SK, Corson K, Hickam DH, Perrin NA, Kraemer DF, Gerrity MS. Depression decision support in primary care - A cluster randomized trial. *Annals of Internal Medicine*. 2006;145:477-487.
- (122) Dobscha SK, Corson K, Pruitt S, Crutchfield M, Gerrity MS. Measuring depression and pain with home health monitors. *Telemedicine Journal and E-Health*. 2006;12:702-706.
- (123) Dobscha SK, Corson K, Gerrity MS. Depression treatment preferences of VA primary care patients. *Psychosomatics*. 2007;48:482-488.

- (124) Dobscha SK, Corson K, Perrin NA et al. Collaborative Care for Chronic Pain in Primary Care A Cluster Randomized Trial. *Jama-Journal of the American Medical Association*. 2009;301:1242-1252.
- (125) Dodds S, Nuehring EM, Blaney NT et al. Integrating mental health services into primary HIV care for women: The Whole Life project. *Public Health Reports*. 2004;119:48-59.
- (126) Domalanta DD, Risser WL, Roberts RE, Risser JMH. Prevalence of depression and other psychiatric disorders among incarcerated youths. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2003;42:477-484.
- (127) Donnelly PL. The use of the patient health questionnaire-9 Korean version (PHQ-9K) to screen for depressive disorders among Korean Americans. *Journal of Transcultural Nursing*. 2007;18:324-330.
- (128) Douglas KM, Taylor AJ, O'Malley PG. Relationship between depression and C-reactive protein in a screening population. *Psychosomatic Medicine*. 2004;66:679-683.
- (129) Drayer RA, Piraino B, Reynolds CF et al. Characteristics of depression in hemodialysis patients: symptoms, quality of life and mortality risk. *General Hospital Psychiatry*. 2006;28:306-312.
- (130) Du Montcel ST, Charles P, Ribai P et al. Composite cerebellar functional severity score: validation of a quantitative score of cerebellar impairment. *Brain*. 2008;131:1352-1361.
- (131) Duffy FF, Chung H, Trivedi M, Rae DS, Regier DA, Katzelnick DJ. Systematic use of patient-rated depression severity monitoring: Is it helpful and feasible in clinical psychiatry? *Psychiatric Services*. 2008;59:1148-1154.
- (132) Dum M, Pickren J, Sobell LC, Sobell MB. Comparing the BDI-II and the PHQ-9 with outpatient substance abusers. *Addictive Behaviors*. 2008;33:381-387.
- (133) Dunkel A, Kendel F, Lehmkuhl E et al. Predictors of preoperative depressive risk in patients undergoing coronary artery bypass graft surgery. *Clinical Research in Cardiology*. 2009;98:643-650.
- (134) Dwight-Johnson M, Ell K, Lee PJ. Can collaborative care address the needs of low-income Latinas with comorbid depression and cancer? Results from a randomized pilot study. *Psychosomatics*. 2005;46:224-232.
- (135) Eack SM, Greeno CG, Lee BJ. Limitations of the Patient Health Questionnaire in identifying anxiety and depression in community mental health: Many cases are undetected. *Research on Social Work Practice*. 2006;16:625-631.
- (136) Eberhard-Gran M, Eskild A, Tambs K, Schel B, Opjordsmoen S. The Edinburgh Postnatal Depression Scale: Validation in a Norwegian community sample. *Nord J Psychiatr*. 2001;55:113-117.
- (137) Eisenman DP, Gelberg L, Liu HH, Shapiro MF. Mental health and health-related quality of life among adult Latino primary care patients living in the United States with previous exposure to political violence. *Jama-Journal of the American Medical Association*. 2003;290:627-634.

- (138) Ekstrand JR, O'Malley PG, Labutta RJ, Jackson JL. The presence of psychiatric disorders reduces the likelihood of neurologic disease among referrals to a neurology clinic. *Journal of Psychosomatic Research*. 2004;57:11-16.
- (139) Ell K, Sanchez K, Vourlekis B et al. Depression, correlates of depression, and receipt of depression care among low-income women with breast or gynecologic cancer. *Journal of Clinical Oncology*. 2005;23:3052-3060.
- (140) Ell K, Quon B, Quinn DI et al. Improving treatment of depression among low-income patients with cancer: the design of the ADAPt-C study. *General Hospital Psychiatry*. 2007;29:223-231.
- (141) Ell K, Xie B, Quon B, Quinn DI, Dwight-Johnson M, Lee PJ. Randomized controlled trial of collaborative care management of depression among low-income patients with cancer. *Journal of Clinical Oncology*. 2008;26:4488-4496.
- (142) Ell K, Xie B, Lee PJ. Impact of Collaborative Care Management of Depression Among Patients With Cancer IN REPLY. *Journal of Clinical Oncology*. 2009;27:1730-1731.
- (143) Ell K, Katon W, Cabassa LJ et al. Depression and Diabetes Among Low-Income Hispanics: Design Elements of A Socio-Culturally Adapted Collaborative Care Model Randomized Controlled Trial. *International Journal of Psychiatry in Medicine*. 2009;39:113-132.
- (144) Esler D, Johnston F, Thomas D, Davis B. The validity of a depression screening tool modified for use with Aboriginal and Torres Strait Islander people. *Australian and New Zealand Journal of Public Health*. 2008;32:317-321.
- (145) Esler DM, Johnston F, Thomas D. The acceptability of a depression screening tool in an urban, Aboriginal community-controlled health service. *Australian and New Zealand Journal of Public Health*. 2007;31:259-263.
- (146) Fan AZ, Strine TW, Jiles R, Mokdad AH. Depression and anxiety associated with cardiovascular disease among persons aged 45 years and older in 38 states of the United States. *Preventive Medicine*. 2008;46:445-450.
- (147) Fann JR, Bombardier CH, Dikmer S et al. Validity of the patient health questionnaire-9 in assessing depression following traumatic brain injury. *Journal of Head Trauma Rehabilitation*. 2005;20:501-511.
- (148) Fann JR, Jones AL, Dikmen SS, Temkin NR, Esselman PC, Bombardier CH. Depression Treatment Preferences After Traumatic Brain Injury. *Journal of Head Trauma Rehabilitation*. 2009;24:272-278.
- (149) Fann JR, Berry DL, Wolpin S et al. Depression screening using the Patient Health Questionnaire-9 administered on a touch screen computer. *Psycho-Oncology*. 2009;18:14-22.
- (150) Ferris DG, Litaker MS, Gilman PA, Lopez AGL. Patient acceptance and the psychological effects of women experiencing telecolposcopy and colposcopy. *Journal of the American Board of Family Practice*. 2003;16:405-411.

- (151) Ferro T, Verdelli H, Pierre F, Weissman MM. Screening for depression in mothers bringing their offspring for evaluation or treatment of depression. *Amer J Psychiat*. 2000;157:375-379.
- (152) Fisher KM, Copenhaver V. Assessing the mental health of rural older adults in public housing facilities: a comparison of screening tools. *J Gerontol Nurs*. 2006;32:26-33.
- (153) Fleishman JA, Zuvekas SH. Global self-rated mental health: Associations with other mental health measures and with role functioning. *Medical Care*. 2007;45:602-609.
- (154) Fogarty CT, Sharma S, Chetty VK, Culpepper L. Mental health conditions are associated with increased health care utilization among urban family medicine patients. *Journal of the American Board of Family Medicine*. 2008;21:398-407.
- (155) Fraguas R, Henriques SG, De Lucia MS et al. The detection of depression in medical setting: A study with PRIME-MD. *Journal of Affective Disorders*. 2006;91:11-17.
- (156) Francis L, Weiss BD, Senf JH, Heist K, Hargraves R. Does literacy education improve symptoms of depression and self-efficacy in individuals with low literacy and depressive symptoms? A preliminary investigation. *Journal of the American Board of Family Medicine*. 2007;20:23-27.
- (157) Freeman EW, Sammel MD, Lin H, Nelson DB. Associations of hormones and menopausal status with depressed mood in women with no history of depression. *Archives of General Psychiatry*. 2006;63:375-382.
- (158) Freynhagen R, Baron R, Tolle T et al. Screening of neuropathic pain components in patients with chronic back pain associated with nerve root compression: a prospective observational pilot study (MIPORT). *Current Medical Research and Opinion*. 2006;22:529-537.
- (159) Gameroff MJ, Olfson M. Major depressive disorder, somatic pain, and health care costs in an urban primary care practice. *Journal of Clinical Psychiatry*. 2006;67:1232-1239.
- (160) Garbers S, Correa N, Tobier N, Blust S, Chiasson MA. Association Between Symptoms of Depression and Contraceptive Method Choices Among Low-Income Women at Urban Reproductive Health Centers. *Maternal and Child Health Journal*. 2010;14:102-109.
- (161) Garcia-Campayo J, Zamorano E, Ruiz MA et al. Cultural adaptation into Spanish of the generalized anxiety disorder-7 (GAD-7) scale as a screening tool. *Health and Quality of Life Outcomes*. 2010;8.
- (162) Garg N, Yates WR, Jones R, Zhou M, Williams S. Effect of gender, treatment site and psychiatric comorbidity on quality of life outcome in substance dependence. *Amer J Addiction*. 1999;8:44-54.
- (163) Garlow SJ, Rosenberg J, Moore JD et al. Depression, desperation, and suicidal ideation in college students: Results from the American Foundation for Suicide Prevention College Screening Project at Emory University. *Depression and Anxiety*. 2008;25:482-488.

- (164) Gary TL, Stark SA, LaVeist TA. Neighborhood characteristics and mental health among African Americans and whites living in a racially integrated urban community. *Health & Place*. 2007;13:569-575.
- (165) Gehi A, Mangano D, Pipkin S, Browner WS, Whooley MA. Depression and heart rate variability in patients with stable coronary heart disease: findings from the Heart and Soul Study. *Arch Gen Psychiatry*. 2005;62:661-666.
- (166) Gehi A, Haas D, Pipkin S, Whooley MA. Depression and medication adherence in outpatients with coronary heart disease: findings from the Heart and Soul Study. *Arch Intern Med*. 2005;165:2508-2513.
- (167) Gehi AK, Ali S, Na B, Whooley MA. Self-reported medication adherence and cardiovascular events in patients with stable coronary heart disease: the heart and soul study. *Arch Intern Med*. 2007;167:1798-1803.
- (168) Gensichen J, Torge M, Peitz M et al. Case management for the treatment of patients with major depression in general practices - rationale, design and conduct of a cluster randomized controlled trial - PRoMPT (Primary care Monitoring for depressive Patient's Trial) [ISRCTN66386086] Study protocol. *Bmc Public Health*. 2005;5:101.
- (169) Gensichen J, Von Korff M, Peitz M et al. Case Management for Depression by Health Care Assistants in Small Primary Care Practices A Cluster Randomized Trial. *Annals of Internal Medicine*. 2009;151:369-W118.
- (170) Gerber MR, Wittenberg E, Ganz ML, Williams CM, McCloskey LA. Intimate partner violence exposure and change in women's physical symptoms over time. *Journal of General Internal Medicine*. 2008;23:64-69.
- (171) Gerrity MS, Corson K, Dobscha SK. Screening for posttraumatic stress disorder in VA primary care patients with depression symptoms. *Journal of General Internal Medicine*. 2007;22:1321-1324.
- (172) Gilbody S, Richards D, Brealey S, Hewitt C. Screening for depression in medical settings with the Patient Health Questionnaire (PHQ): a diagnostic meta-analysis. *J Gen Intern Med*. 2007;22:1596-1602.
- (173) Gilbody S, Richards D, Barkham M. Diagnosing depression in primary care using self-completed instruments: UK validation of PHQ-9 and CORE-OM. *British Journal of General Practice*. 2007;57:650-652.
- (174) Gilmer TP, Walker C, Johnson ED, Philis-Tsimikas A, Unutzer J. Improving treatment of depression among Latinos with diabetes using Project Dulce and IMPACT. *Diabetes Care*. 2008;31:1324-1326.
- (175) Gjerdingen D, Crow S, McGovern P, Miner M, Center B. Postpartum Depression Screening at Well-Child Visits: Validity of a 2-Question Screen and the PHQ-9. *Annals of Family Medicine*. 2009;7:63-70.

- (176) Glaesmer H, Gunzelmann T, Martin A, Brahler E, Rief W. The impact of mental disorders on health care utilization and illness behaviour in the elderly. *Psychiatrische Praxis*. 2008;35:187-193.
- (177) Glaesmer H, Wittig U, Brahler E, Martin A, Mewes R, Rief W. Are Migrants More Susceptible to Mental Disorders? An Evaluation of a Representative Sample of the German General Population. *Psychiatrische Praxis*. 2009;36:16-22.
- (178) Glasgow RE, Nutting PA, King DK et al. A practical randomized trial to improve diabetes care. *Journal of General Internal Medicine*. 2004;19:1167-1174.
- (179) Glasgow RE, Nutting PA, King DK et al. Randomized effectiveness trial of a computer-assisted intervention to improve diabetes care. *Diabetes Care*. 2005;28:33-39.
- (180) Glass RM. Minds and machines. Interactive voice response technology to detect mental disorders. *JAMA*. 1997;278:945-946.
- (181) Goebel JR, Doering LV, Shugarman LR et al. Heart Failure: The Hidden Problem of Pain. *Journal of Pain and Symptom Management*. 2009;38:698-707.
- (182) Goebel S, Baumann B, Steinert A, Reppenhagen S, Faller H. Elevated postoperative pain levels following orthopedic surgery. Depression as a strong predictor. *Schmerz*. 2010;24:54-61.
- (183) Goebert D, Thompson D, Takeshita J et al. Depressive Symptoms in Medical Students and Residents: A Multischool Study. *Academic Medicine*. 2009;84:236-241.
- (184) Goldney R, Hawthorne G, Fisher L. Is the Australian National Survey of Mental Health and Wellbeing a reliable guide for health planners? A methodological note on the prevalence of depression. *Australian and New Zealand Journal of Psychiatry*. 2004;38:635-638.
- (185) Goldney RD, Fisher LJ, Wilson DH. Mental health literacy: an impediment to the optimum treatment of major depression in the community. *J Affect Disorders*. 2001;64:277-284.
- (186) Goldney RD, Ruffin R, Fisher LJ, Wilson DH. Asthma symptoms associated with depression and lower quality of life: a population survey. *Medical Journal of Australia*. 2003;178:437-441.
- (187) Goldney RD, Fisher LJ. Double depression in an Australian population. *Social Psychiatry and Psychiatric Epidemiology*. 2004;39:921-926.
- (188) Goldney RD, Fisher LJ, Dal Grande E, Taylor AW. Subsyndromal depression: prevalence, use of health services and quality of life in an Australian population. *Social Psychiatry and Psychiatric Epidemiology*. 2004;39:293-298.
- (189) Goldney RD, Bain MA. Double depression: its morbidity and management in a community setting. *Acta Neuropsychiatrica*. 2006;18:210-215.
- (190) Goldney RD, Fisher LJ, Dal Grande E, Taylor AW, Hawthorne G. Have education and publicity about depression made a difference? Comparison of prevalence, service use and excess costs in South Australia: 1998 and 2004. *Australian and New Zealand Journal of Psychiatry*. 2007;41:38-53.

- (191) Goldney RD, Dunn KI, Air TM, Dal Grande E, Taylor AW. Relationships between body mass index, mental health, and suicidal ideation: population perspective using two methods. *Australian and New Zealand Journal of Psychiatry*. 2009;43:652-658.
- (192) Goodwin R, Olfson M, Feder A, Fuentes M, Pilowsky DJ, Weissman MM. Panic and suicidal ideation in primary care. *Depress Anxiety*. 2001;14:244-246.
- (193) Goodwin R, Olfson M. Self-perception of poor health and suicidal ideation in medical patients. *Psychological Medicine*. 2002;32:1293-1299.
- (194) Goodwin RD, Olfson M, Shea S et al. Asthma and mental disorders in primary care. *General Hospital Psychiatry*. 2003;25:479-483.
- (195) Goodwin RD, Hoven CW, Spitzer RL. Diabetes and eating disorders in primary care. *International Journal of Eating Disorders*. 2003;33:85-91.
- (196) Goodwin RD, Kroenke K, Hoven CW, Spitzer RL. Major depression, physical illness, and suicidal ideation in primary care. *Psychosomatic Medicine*. 2003;65:501-505.
- (197) Grafe K, Zipfel S, Herzog W, Lowe B. Screening for psychiatric disorders with the Patient Health Questionnaire (PHQ). Results from the German validation study. *Diagnostica*. 2004;50:171-181.
- (198) Grandy S, Chapman RH, Fox KM. Quality of life and depression of people living with type 2 diabetes mellitus and those at low and high risk for type 2 diabetes: findings from the Study to Help Improve Early evaluation and management of risk factors Leading to Diabetes (SHIELD). *International Journal of Clinical Practice*. 2008;62:562-568.
- (199) Graves DE, Bombardier CH. Improving the efficiency of screening for major depression in people with spinal cord injury. *Journal of Spinal Cord Medicine*. 2008;31:177-184.
- (200) Gray GC, Chesbrough KB, Ryan MAK et al. The millennium cohort study: A 21-year prospective cohort study, of 140,000 military personnel. *Military Medicine*. 2002;167:483-488.
- (201) Green MS, Naumann RW, Elliot M, Hall JB, Higgins RV, Grigsby JH. Sexual dysfunction following vulvectomy. *Gynecol Oncol*. 2000;77:73-77.
- (202) Greiver M, Anderson E, van Weel-Baumgarten E. Another approach to managing depression. *Canadian Family Physician*. 2005;51:25.
- (203) Grieger TA, Waldrep DA, Lovasz MM, Ursano RJ. Follow-up of Pentagon employees two years after the terrorist attack of September 11, 2001. *Psychiatric Services*. 2005;56:1374-1378.
- (204) Grieger TA, Cozza SJ, Ursano RJ et al. Posttraumatic stress disorder and depression in battle-injured soldiers. *American Journal of Psychiatry*. 2006;163:1777-1783.
- (205) Grucza RA, Przybeck TR, Cloninger CR. Prevalence and correlates of binge eating disorder in a community sample. *Comprehensive Psychiatry*. 2007;48:124-131.

- (206) Grupp-Phelan J, Whitaker RC, Naish AB. Depression in mothers of children presenting for emergency and primary care: Impact on mothers' perceptions of caring for their children. *Ambulatory Pediatrics*. 2003;3:142-146.
- (207) Grypma L, Haverkamp R, Little S, Unutzer J. Taking an evidence-based model of depression care from research to practice: making lemonade out of depression. *General Hospital Psychiatry*. 2006;28:101-107.
- (208) Guimaraes JMN, Lopes CD, Baima J, Sichieri R. Depression symptoms and hypothyroidism in a population-based study of middle-aged Brazilian women. *Journal of Affective Disorders*. 2009;117:120-123.
- (209) Gunn JM, Gilchrist GP, Chondros P et al. Who is identified when screening for depression is undertaken in general practice? Baseline findings from the Diagnosis, Management and Outcomes of Depression in Primary Care (diamond) longitudinal study. *Medical Journal of Australia*. 2008;188:S119-S125.
- (210) Gupta MX, Silberstein SD, Young WB, Hopkins M, Lopez BL, Samsa GP. Less is not more: Underutilization of headache medications in a university hospital emergency department. *Headache*. 2007;47:1125-1133.
- (211) Haas A, Koestner B, Rosenberg J et al. An interactive Web-based method of outreach to college students at risk for suicide. *Journal of American College Health*. 2008;57:15-22.
- (212) Haas AP, Hendin H, Mann JJ. Suicide in college students. *American Behavioral Scientist*. 2003;46:1224-1240.
- (213) Haggarty J, O'Connor B, Dubois S et al. A pilot study of a Canadian shared mental health care programme: Changes in patient symptoms and disability. *Primary Care & Community Psychiatry*. 2008;13:27-35.
- (214) Haggman S, Maher CG, Refshauge KM. Screening for symptoms of depression by physical therapists managing low back pain. *Physical Therapy*. 2004;84:1157-1166.
- (215) Hahn SR, Kroenke K, Spitzer RL et al. The difficult patient: Prevalence, psychopathology, and functional impairment. *J Gen Intern Med*. 1996;11:1-8.
- (216) Hahn SR. Physical symptoms and physician-experienced difficulty in the physician-patient relationship. *Ann Intern Med*. 2001;134:897-904.
- (217) Han C, Pae CU, Lee BH et al. Fluoxetine versus sertraline in the treatment of patients with undifferentiated somatoform disorder: A randomized, open-label, 12-week, parallel-group trial. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 2008;32:437-444.
- (218) Han C, Pae CU, Lee BH et al. Venlafaxine versus mirtazapine in the treatment of undifferentiated somatoform disorder - A 12-week prospective, open-label, randomized, parallel-group trial. *Clinical Drug Investigation*. 2008;28:251-261.

- (219) Han C, Pae CU, Patkar AA et al. Psychometric Properties of the Patient Health Questionnaire-15 (PHQ-15) for Measuring the Somatic Symptoms of Psychiatric Outpatients. *Psychosomatics*. 2009;50:580-585.
- (220) Han CS, Marks DM, Pae CU et al. Paroxetine for patients with undifferentiated somatoform disorder: A prospective, open-label, 8-week pilot study. *Current Therapeutic Research-Clinical and Experimental*. 2008;69:221-231.
- (221) Han CS, Jo SA, Kwak JH et al. Validation of the Patient Health Questionnaire-9 Korean version in the elderly population: the Ansan Geriatric Study. *Comprehensive Psychiatry*. 2008;49:218-223.
- (222) Hancock P, Lerner AJ. Clinical utility of Patient Health Questionnaire-9 (PHQ-9) in memory clinics. *International Journal of Psychiatry in Clinical Practice*. 2009;13:188-191.
- (223) Hanel G, Henningsen P, Herzog W et al. Depression, anxiety, and somatoform disorders: Vague or distinct categories in primary care? Results from a large cross-sectional study. *Journal of Psychosomatic Research*. 2009;67:189-197.
- (224) Hansson M, Chotai J, Nordstrom A, Bodlund O. Comparison of two self-rating scales to detect depression: HADS and PHQ-9. *British Journal of General Practice*. 2009;59:650-654.
- (225) Hanusa BH, Scholle SH, Haskett RF, Spadaro K, Wisner KL. Screening for depression in the postpartum period: A comparison of three instruments. *Journal of Womens Health*. 2008;17:585-596.
- (226) Hauser W, Schmutzer G, Glaesmer H, Brahler E. Prevalence and predictors of pain in several body regions - Results of a representative German population survey. *Schmerz*. 2009;23:461-+.
- (227) Hauser W, Schmutzer G, Brahler E, Glaesmer H. A Cluster Within the Continuum of Biopsychosocial Distress Can Be Labeled "Fibromyalgia Syndrome" - Evidence from a Representative German Population Survey. *Journal of Rheumatology*. 2009;36:2806-2812.
- (228) Haut SR, Katz M, Masur J, Lipton RB. Seizures in the elderly: Impact on mental status, mood, and sleep. *Epilepsy & Behavior*. 2009;14:540-544.
- (229) Hawthorne G, Cheek F, Goldney R, Fisher L. The excess cost of depression in South Australia: a population-based study. *Australian and New Zealand Journal of Psychiatry*. 2003;37:362-373.
- (230) Hawthorne G, Goldney R, Taylor AW. Depression prevalence: is it really increasing? *Australian and New Zealand Journal of Psychiatry*. 2008;42:606-616.
- (231) Hegel MT, Collins ED, Kearing S, Gillock KL, Moore CP, Ahles TA. Sensitivity and specificity of the Distress Thermometer for depression in newly diagnosed breast cancer patients. *Psycho-Oncology*. 2008;17:556-560.
- (232) Henkel V, Mergl R, Kohnen R, Maier W, Moller HJ, Hegerl U. Identifying depression in primary care: a comparison of different methods in a prospective cohort study. *British Medical Journal*. 2003;326:200-201.

- (233) Henkel V, Mergl R, Coyne JC, Kohnen R, Moller HJ, Hegerl U. Screening for depression in primary care: Will one or two items suffice? *European Archives of Psychiatry and Clinical Neuroscience*. 2004;254:215-223.
- (234) Henkel V, Mergl R, Kohnen R, Allgaier AK, Moller HJ, Hegerl U. Use of brief depression screening tools in primary care: consideration of heterogeneity in performance in different patient groups. *General Hospital Psychiatry*. 2004;26:190-198.
- (235) Henriques SG, Fraguas R, Iosifescu DV et al. Recognition of Depressive Symptoms by Physicians. *Clinics*. 2009;64:629-635.
- (236) Hepner KA, Morales LS, Hays RD, Edelen MO, Miranda J. Evaluating differential item functioning of the PRIME-MD mood module among impoverished black and white women in primary care. *Womens Health Issues*. 2008;18:53-61.
- (237) Hepner KA, Hunter SB, Edelen MO, Zhou AJ, Watkins K. A comparison of two depressive symptomatology measures in residential substance abuse treatment clients. *Journal of Substance Abuse Treatment*. 2009;37:318-325.
- (238) Hepner KA, Hunter SB, Edelen MO, Zhou AJ, Watkins K. A comparison of two depressive symptomatology measures in residential substance abuse treatment clients. *J Subst Abuse Treat*. 2009.
- (239) Hides L, Lubman DI, Devlin H et al. Reliability and validity of the Kessler 10 and Patient Health Questionnaire among injecting drug users. *Australian and New Zealand Journal of Psychiatry*. 2007;41:166-168.
- (240) Hirshfield S, Wolitski RJ, Chiasson A, Remien RH, Humberstone M, Wong T. Screening for depressive symptoms in an online sample of men who have sex with men. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2008;20:904-910.
- (241) Hoedeman R, Krol B, Blankenstein N, Koopmans PC, Groothoff JW. Severe MUPS in a sick-listed population: a cross-sectional study on prevalence, recognition, psychiatric co-morbidity and impairment. *Bmc Public Health*. 2009;9.
- (242) Hoge CW, McGurk D, Thomas JL, Cox AL, Engel CC, Castro CA. Mild traumatic brain injury in U.S. Soldiers returning from Iraq. *N Engl J Med*. 2008;358:453-463.
- (243) Hollinrake E, Abreu A, Maifeld M, Van Voorhis BJ, Dokras A. Increased risk of depressive disorders in women with polycystic ovary syndrome. *Fertil Steril*. 2007;87:1369-1376.
- (244) Holzapfel N, Zugck C, Muller-Tasch T et al. Routine screening for depression and quality of life in outpatients with congestive heart failure. *Psychosomatics*. 2007;48:112-116.
- (245) Holzapfel N, Muffler-Tasch T, Wild B et al. Depression profile in patients with and without chronic heart failure. *Journal of Affective Disorders*. 2008;105:53-62.
- (246) Horn EK, van Benthem TB, Roijen LHV et al. Cost-effectiveness of collaborative care for chronically ill patients with comorbid depressive disorder in the general hospital setting, a randomised controlled trial. *Bmc Health Services Research*. 2007;7.

- (247) Howell EA, Mora PA, DiBonaventura MD, Leventhal H. Modifiable factors associated with changes in postpartum depressive symptoms. *Archives of Womens Mental Health*. 2009;12:113-120.
- (248) Huang FY, Chung H, Kroenke K, Delucchi KL, Spitzer RL. Using the patient health questionnaire-9 to measure depression among racially and ethnically diverse primary care patients. *Journal of General Internal Medicine*. 2006;21:547-552.
- (249) Huang FY, Chung H, Kroenke K, Spitzer RL. Racial and ethnic differences in the relationship between depression severity and functional status. *Psychiatric Services*. 2006;57:498-503.
- (250) Hunter CL, Hunter CM, West ET, Kinder MH, Carroll DW. Recognition of depressive disorders by primary care providers in a military medical setting. *Military Medicine*. 2002;167:308-311.
- (251) Husain N, Creed F, Tomenson B. Depression and social stress in Pakistan. *Psychol Med*. 2000;30:395-402.
- (252) Husain N, Waheed W, Tomenson B, Creed F. The validation of personal health questionnaire amongst people of Pakistani family origin living in the United Kingdom. *Journal of Affective Disorders*. 2007;97:261-264.
- (253) Interian A, Allen LA, Gara MA, Escobar JI, az-Martinez AM. Somatic complaints in primary care: Further examining the validity of the patient health questionnaire (PHQ-15). *Psychosomatics*. 2006;47:392-398.
- (254) Interian A, Ang A, Gara MA, Link BG, Rodriguez MA, Vega WA. Stigma and Depression Treatment Utilization Among Latinos: Utility of Four Stigma Measures. *Psychiatric Services*. 2010;61:373-379.
- (255) Iversen AC, van Staden L, Hughes JH et al. The prevalence of common mental disorders and PTSD in the UK military: using data from a clinical interview-based study. *Bmc Psychiatry*. 2009;9.
- (256) Jackson JL, Kroenke K. Difficult patient encounters in the ambulatory clinic - Clinical predictors and outcomes. *Arch Intern Med*. 1999;159:1069-1075.
- (257) Jackson JL, O'Malley PG, Kroenke K. A psychometric comparison of military and civilian medical practices. *Milit Med*. 1999;164:112-115.
- (258) Jackson JL, Kroenke K, Chamberlin J. Effects of physician awareness of symptom-related expectations and mental disorders. A controlled trial. *Arch Fam Med*. 1999;8:135-142.
- (259) Jackson JL, Houston JS, Hanling SR, Terhaar KA, Yun JS. Clinical predictors of mental disorders among medical outpatients. *Arch Intern Med*. 2001;161:875-879.
- (260) Jackson JL, Kroenke K. The effect of unmet expectations among adults presenting with physical symptoms. *Ann Intern Med*. 2001;134:889-897.
- (261) Jackson JL, Chamberlin J, Kroenke K. Predictors of patient satisfaction. *Soc Sci Med*. 2001;52:609-620.

- (262) Jackson JL. Communication about symptoms in primary care: Impact on patient outcomes. *Journal of Alternative and Complementary Medicine*. 2005;11:S51-S56.
- (263) Jackson JL, Passamonti M, Kroenke K. Outcome and impact of mental disorders in primary care at 5 years. *Psychosom Med*. 2007;69:270-276.
- (264) Jacobs SR, Jacobsen PB, Donovan K, Booth-Jones M. Utility of the Patient Health Questionnaire-9 (Phq-9) in Identifying Depression Among Hematopoietic Stem Cell Transplant (Hsct) Patients. *Annals of Behavioral Medicine*. 2007;33:S56.
- (265) Jacobson IG, Ryan MA, Hooper TI et al. Alcohol use and alcohol-related problems before and after military combat deployment. *JAMA*. 2008;300:663-675.
- (266) Jacobson IG, White MR, Smith TC et al. Self-Reported Health Symptoms and Conditions Among Complementary and Alternative Medicine Users in a Large Military Cohort. *Annals of Epidemiology*. 2009;19:613-622.
- (267) Jacobson IG, Smith TC, Smith B et al. Disordered eating and weight changes after deployment: longitudinal assessment of a large US military cohort. *Am J Epidemiol*. 2009;169:415-427.
- (268) Jarjoura D, Polen A, Baum E, Kropp D, Hetrick S, Rutecki G. Effectiveness of screening and treatment for depression in ambulatory indigent patients. *Journal of General Internal Medicine*. 2004;19:78-84.
- (269) Jefford M, Mileskin L, Richards K et al. Rapid screening for depression - validation of the Brief Case-Find for Depression (BCD) in medical oncology and palliative care patients. *British Journal of Cancer*. 2004;91:900-906.
- (270) Jeon WS, Park TJ, Kim SY, Pode R, Jang J, Kwon JH. Ideal host and guest system in phosphorescent OLEDs. *Organic Electronics*. 2009;10:240-246.
- (271) Jewell JS, Dunn AL, Bondy J, Leiferman J. Prevalence of Self-Reported Postpartum Depression Specific to Season and Latitude of Birth: Evaluating the PRAMS Data. *Maternal and Child Health Journal*. 2010;14:261-267.
- (272) Johnson GD, Formichella C, Thomas JS, Bhaumik D, Degruy F, Riordan CA. Stress and distress among Gulf of Mexico shrimp fishermen. *Hum Organ*. 1998;57:404-413.
- (273) JOHNSON JG, Spitzer RL, Kroenke K et al. Psychiatric comorbidity, health-status, and functional impairment associated with alcohol-abuse and dependence in primary-care patients - Findings of the PRIME-MD 1000 Study. *J Consult Clin Psychol*. 1995;63:133-140.
- (274) JOHNSON JG, Harris ES, Spitzer RL, Williams JBW. The Patient Health Questionnaire for Adolescents: Validation of an instrument for the assessment of mental disorders among adolescent primary care patients. *J Adolescent Health*. 2002;30:196-204.
- (275) JOHNSON JG, Spitzer RS, Kroenke K, Williams JBW. *Manual and scoring instructions for the Patient Health Questionnaire for Adolescents (PHQ-A)*. New York, NY: Biometrics Research Department, New York State Psychiatric Institute; 2005.

- (276) JOHNSON JG, Spitzer RS, Kroenke K, Williams JBW. *The Patient Health Questionnaire for Adolescents, Revised Version (PHQ-A)*. New York, NY: Biometrics Research Department, New York State Psychiatric Institute; 2005.
- (277) Johnson RL, Botwinick G, Sell RL et al. The utilization of treatment and case management services by HIV-infected youth. *Journal of Adolescent Health*. 2003;33:31-38.
- (278) Jonkers CCM, Lamers F, Evers SMAA, Bosma H, Metsemakers JF, van Eijk JTM. Economic evaluation of a minimal psychological intervention in chronically ill elderly patients with minor or mild to moderate depression.: A randomized trial (the DELTA-study). *International Journal of Technology Assessment in Health Care*. 2009;25:497-504.
- (279) Joubert J, Joubert L, Reid C et al. The positive effect of integrated care on depressive symptoms in stroke survivors. *Cerebrovascular Diseases*. 2008;26:199-205.
- (280) Justice AC, McGinnis KA, Atkinson JH et al. Psychiatric and neurocognitive disorders among HIV-positive and negative veterans in care: Veterans Aging Cohort Five-Site Study. *Aids*. 2004;18:S49-S59.
- (281) Kahn LS, Fox CH, McIntyre RS, Tumiel-Berhalter L, Berdine DE, Lyle H. Assessing the prevalence of depression among individuals with diabetes in a Medicaid managed-care program. *International Journal of Psychiatry in Medicine*. 2008;38:13-29.
- (282) Kalpakjian CZ, Albright KJ. An examination of depression through the lens of spinal cord injury - Comparative prevalence rates and severity in women and men. *Womens Health Issues*. 2006;16:380-388.
- (283) Kalpakjian CZ, Toussaint LL, Albright KJ, Bombardier CH, Krause JK, Tate DG. Patient Health Questionnaire-9 in Spinal Cord Injury: An Examination of Factor Structure as Related to Gender. *Journal of Spinal Cord Medicine*. 2009;32:147-156.
- (284) Katon W, Von Korff M, Ciechanowski P et al. Behavioral and clinical factors associated with depression among individuals with diabetes. *Diabetes Care*. 2004;27:914-920.
- (285) Katon W, Unutzer J, Russo J. Major depression: the importance of clinical characteristics and treatment response to prognosis. *Depress Anxiety*. 2009.
- (286) Katon W, Russo J, Lin EHB et al. Diabetes and Poor Disease Control: Is Comorbid Depression Associated With Poor Medication Adherence or Lack of Treatment Intensification? *Psychosomatic Medicine*. 2009;71:965-972.
- (287) Katon WJ, Lin E, Russo J, Unutzer J. Increased medical costs of a population-based sample of depressed elderly patients. *Archives of General Psychiatry*. 2003;60:897-903.
- (288) Katon WJ, Simon G, Russo J et al. Quality of depression care in a population-based sample of patients with diabetes and major depression. *Medical Care*. 2004;42:1222-1229.
- (289) Katon WJ. Panic disorder. *New England Journal of Medicine*. 2006;354:2360-2U20.

- (290) Kelly R, Zatzick D, Anders T. The detection and treatment of psychiatric disorders and substance use among pregnant women cared for in obstetrics. *Am J Psychiatry*. 2001;158:213-219.
- (291) Kelly RH, Russo J, Katon W. Somatic complaints among pregnant women cared for in obstetrics: normal pregnancy or depressive and anxiety symptom amplification revisited? *Gen Hosp Psychiat*. 2001;23:107-113.
- (292) Kendrick T, Dowrick C, McBride A et al. Management of depression in UK general practice in relation to scores on depression severity questionnaires: analysis of medical record data. *British Medical Journal*. 2009;338.
- (293) Kerchner A, Lester W, Stuart SP, Dokras A. Risk of depression and other mental health disorders in women with polycystic ovary syndrome: a longitudinal study. *Fertility and Sterility*. 2009;91:207-212.
- (294) Kerse N, Flicker L, Pfaff JJ et al. Falls, Depression and Antidepressants in Later Life: A Large Primary Care Appraisal. *Plos One*. 2008;3.
- (295) Ketterer MW, Wulsin L, Cao JJ et al. "Major" depressive disorder, coronary heart disease, and the DSM-IV threshold problem. *Psychosomatics*. 2006;47:50-55.
- (296) Ketterer MW, Knysz W. Screening, diagnosis & monitoring of depression/distress in CHF patients. *Heart Failure Reviews*. 2009;14:1-5.
- (297) Kilbourne AM, Reynolds CF, Good CB, Sereika SM, Justice AC, Fine MJ. How does depression influence diabetes medication adherence in older patients? *American Journal of Geriatric Psychiatry*. 2005;13:202-210.
- (298) Kim HG, Mandell M, Crandall C, Kuskowski MA, Dieperink B, Buchberger RL. Antenatal psychiatric illness and adequacy of prenatal care in an ethnically diverse inner-city obstetric population. *Archives of Womens Mental Health*. 2006;9:103-107.
- (299) Kissane DW, Love A, Lee XQ, Kee PL. The demoralization scale: A report of its development and preliminary validation. *Journal of Palliative Care*. 2004;20:269-276.
- (300) Klapow J, Kroenke K, Horton T, Schmidt S, Spitzer R, Williams JBW. Psychological disorders and distress in older primary care patients: A comparison of older and younger samples. *Psychosom Med*. 2002;64:635-643.
- (301) Kobak KA, Taylor LV, Dottl SL et al. A computer-administered telephone interview to identify mental disorders. *JAMA*. 1997;278:905-910.
- (302) Kobak KA, Taylor LV, Dottl SL et al. Computerized screening for psychiatric disorders in an outpatient community mental health clinic. *Psychiatr Services*. 1997;48:1048-1057.
- (303) Kobrosly R, van Wijngaarden E. Associations between immunologic, inflammatory, and oxidative stress markers with severity of depressive symptoms: An analysis of the 2005-2006 National Health and Nutrition Examination Survey. *Neurotoxicology*. 2010;31:126-133.

- (304) Kochhar PH, Rajadhyaksha SS, Suvarna VR. Translation and validation of brief patient health questionnaire against DSM IV as a tool to diagnose major depressive disorder in Indian patients. *Journal of Postgraduate Medicine*. 2007;53:102-107.
- (305) Kovac SH, Klapow JC, Kroenke K, Spitzer RL, Williams JBW. Differing symptoms of abused versus nonabused women in obstetric-gynecology settings. *American Journal of Obstetrics and Gynecology*. 2003;188:707-713.
- (306) Kraemer B, Moergeli H, Roth H, Hepp U, Schnyder U. Contribution of initial heart rate to the prediction of posttraumatic stress symptom level in accident victims. *Journal of Psychiatric Research*. 2008;42:158-162.
- (307) Krause JS, Bombardier C, Carter RE. Assessment of Depressive Symptoms During Inpatient Rehabilitation for Spinal Cord Injury: Is There an Underlying Somatic Factor When Using the PHQ? *Rehabilitation Psychology*. 2008;53:513-520.
- (308) Krause JS, Saunders LL, Reed KS, Coker J, Zhai YS, Johnson E. Comparison of the Patient Health Questionnaire and the Older Adult Health and Mood Questionnaire for Self-Reported Depressive Symptoms After Spinal Cord Injury. *Rehabilitation Psychology*. 2009;54:440-448.
- (309) Krause JS, Reed KS, McArdle JJ. Prediction of somatic and non-somatic depressive symptoms between inpatient rehabilitation and follow-up. *Spinal Cord*. 2010;48:239-244.
- (310) Krebs EE, Bair MJ, Carey TS, Weinberger M. Documentation of Pain Care Processes Does Not Accurately Reflect Pain Management Delivered in Primary Care. *Journal of General Internal Medicine*. 2010;25:194-199.
- (311) Krefetz DG, Steer RA, Jermyn RT, Condoluci DV. Screening HIV-infected patients with chronic pain for anxiety and mood disorders with the Beck Anxiety and Depression Inventory-Fast Screens for Medical Settings. *Journal of Clinical Psychology in Medical Settings*. 2004;11:283-289.
- (312) Kroenke K, Spitzer RL, Williams JB et al. Physical symptoms in primary care. Predictors of psychiatric disorders and functional impairment. *Arch Fam Med*. 1994;3:774-779.
- (313) Kroenke K, Jackson JL, Chamberlin J. Depressive and anxiety disorders in patients presenting with physical complaints: Clinical predictors and outcome. *Amer J Med*. 1997;103:339-347.
- (314) Kroenke K, Spitzer RL, deGruy FV, III et al. Multisomatoform disorder. An alternative to undifferentiated somatoform disorder for the somatizing patient in primary care. *Arch Gen Psychiatry*. 1997;54:352-358.
- (315) Kroenke K, Spitzer RL, deGruy FV, III, Swindle R. A symptom checklist to screen for somatoform disorders in primary care. *Psychosomatics*. 1998;39:263-272.
- (316) Kroenke K, Spitzer RL. Gender differences in the reporting of physical and somatoform symptoms. *Psychosom Med*. 1998;60:150-155.
- (317) Kroenke K, Spitzer RL, Williams JBW. The PHQ-9 - Validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16:606-613.

- (318) Kroenke K, Spitzer RL, Williams JBW. The PHQ-15: Validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med*. 2002;64:258-266.
- (319) Kroenke K, Spitzer RL. The PHQ-9: A new depression diagnostic and severity measure. *Psychiatric Annals*. 2002;32:509-515.
- (320) Kroenke K, Spitzer RL, Williams JBW. The Patient Health Questionnaire-2 - Validity of a two-item depression screener. *Medical Care*. 2003;41:1284-1292.
- (321) Kroenke K, Messina N, Benattia I, Graepel J, Musgnung J. Venlafaxine extended release in the short-term treatment of depressed and anxious primary care patients with multisomatoform disorder. *Journal of Clinical Psychiatry*. 2006;67:72-80.
- (322) Kroenke K. Physical symptom disorder: A simpler diagnostic category for somatization-spectrum conditions. *Journal of Psychosomatic Research*. 2006;60:335-339.
- (323) Kroenke K, Bair M, Damush T et al. Stepped Care for Affective Disorders and Musculoskeletal Pain (SCAMP) study design and practical implications of an intervention for comorbid pain and depression. *General Hospital Psychiatry*. 2007;29:506-517.
- (324) Kroenke K, Spitzer RL, Williams JBW, Monahan PO, Lowe B. Anxiety disorders in primary care: Prevalence, impairment, comorbidity, and detection. *Annals of Internal Medicine*. 2007;146:317-325.
- (325) Kroenke K, Spitzer RL, Williams JBW, Lowe B. An Ultra-Brief Screening Scale for Anxiety and Depression: The PHQ-4. *Psychosomatics*. 2009;50:613-621.
- (326) Kroenke K, Strine TW, Spitzer RL, Williams JBW, Berry JT, Mokdad AH. The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*. 2009;114:163-173.
- (327) Kubiak SP, Beeble ML, Bybee D. Testing the Validity of the K6 in Detecting Major Depression and Ptsd Among Jailed Women. *Criminal Justice and Behavior*. 2010;37:64-80.
- (328) Kuehn BM. Scientists examine primary care-based screening and treatments for anxiety. *Jama-Journal of the American Medical Association*. 2008;299:1886-1887.
- (329) Kumar G, Kim AH, Krefetz D, Steer RA. Screening for major depressive disorders in adolescent psychiatric inpatients with the mood modules from the primary care evaluation of mental disorders and the patient health questionnaire. *Psychol Rep*. 2001;89:274-278.
- (330) Kumar G, Steer RA, Teitelman KB, Villacis L. Effectiveness of Beck Depression Inventory-II subscales in screening for major depressive disorders in adolescent psychiatric inpatients. *Assessment*. 2002;9:164-170.
- (331) Kunik ME, Roundy K, Veazey C et al. Surprisingly high prevalence of anxiety and depression in chronic breathing disorders. *Chest*. 2005;127:1205-1211.
- (332) Kunik ME, Azzam PN, Souchek J et al. A practical screening tool for anxiety and depression in patients with chronic breathing disorders. *Psychosomatics*. 2007;48:16-21.

- (333) Kuppermann M, Learman LA, Schembri M et al. Effect of noncancerous pelvic problems on health-related quality of life and sexual functioning. *Obstetrics and Gynecology*. 2007;110:633-642.
- (334) Kurland JE, Coyle WJ, Winkler A, Zable E. Prevalence of irritable bowel syndrome and depression in fibromyalgia. *Digestive Diseases and Sciences*. 2006;51:454-460.
- (335) Kurz B, Malcolm B, Cournoyer D. In the shadow of race: Immigrant status and mental health. *Affilia-Journal of Women and Social Work*. 2005;20:434-447.
- (336) Kuwert P, Brahler E, Glaesmer H, Freyberger HJ, Decker O. Impact of forced displacement during World War II on the present-day mental health of the elderly: a population-based study. *International Psychogeriatrics*. 2009;21:748-753.
- (337) Kuyken W, Watkins E, Holden E, Cook W. Rumination in adolescents at risk for depression. *J Affect Disord*. 2006;96:39-47.
- (338) Lamers F, Jonkers CCM, Bosma H, Penninx BWJH, Knottnerus JA, van Eijk JTM. Summed score of the Patient Health Questionnaire-9 was a reliable and valid method for depression screening in chronically ill elderly patients. *Journal of Clinical Epidemiology*. 2008;61:679-687.
- (339) Lamoureux EL, Tee HW, Pesudovs K, Pallant JF, Keeffe JE, Rees G. Can Clinicians Use the PHQ-9 to Assess Depression in People with Vision Loss? *Optometry and Vision Science*. 2009;86:139-145.
- (340) Lasprilla JCA, Moreno A, Rogers H, Francis K. The Effect of Dementia Patient's Physical, Cognitive, and Emotional/Behavioral Problems on Caregiver Well-Being: Findings From a Spanish-Speaking Sample From Colombia, South America. *American Journal of Alzheimers Disease and Other Dementias*. 2009;24:384-395.
- (341) Lasser KE, Zeytinoglu H, Miller E, Becker AE, Hermann RC, Bor DH. Do women who screen positive for mental disorders in primary care have lower mammography rates? *General Hospital Psychiatry*. 2003;25:214-216.
- (342) Lavan H, JOHNSON JG. The association between Axis I and II psychiatric symptoms and high-risk sexual behavior during adolescence. *Journal of Personality Disorders*. 2002;16:73-94.
- (343) Lavoie KL, Fleet RP, Lesperance F et al. Are exercise stress tests appropriate for assessing myocardial ischemia in patients with major depressive disorder? *American Heart Journal*. 2004;148:621-627.
- (344) Lee KS, Lennie TA, Heo S, Wu JR, Chung ML, Moser DK. Comparison of the Impact of PHQ-9 Scores with and without Physical Symptom Items on Event-Free Survival in Patients with Heart Failure. *Journal of Cardiac Failure*. 2009;15:011.
- (345) Lee PW, Schulberg HC, Raue PJ, Kroenke K. Concordance between the PHQ-9 and the HSCL-20 in depressed primary care patients. *Journal of Affective Disorders*. 2007;99:139-145.
- (346) Lefevre F, Reifler D, Lee P et al. Screening for undetected mental disorders in high utilizers of primary care services. *J Gen Intern Med*. 1999;14:425-431.

- (347) Leithner K, ssem-Hilger E, Fischer-Kern M, Loeffler-Stastka H, Sam C, Ponocny-Seliger E. Psychiatric morbidity in gynecological and otorhinolaryngological outpatients: a comparative study. *General Hospital Psychiatry*. 2009;31:233-239.
- (348) Lenze EJ, Munin MC, Ferrell RE et al. Association of the serotonin transporter gene-linked polymorphic region (5-HTTLPR) genotype with depression in elderly persons after hip fracture. *American Journal of Geriatric Psychiatry*. 2005;13:428-432.
- (349) Leopold KA, Ahles TA, Walch S et al. Prevalence of mood disorders and utility of the PRIME-MD in patients undergoing radiation therapy. *Int J Radiat Oncol Biol Phys*. 1998;42:1105-1112.
- (350) Li CY, Friedman B, Conwell Y, Fiscella K. Validity of the Patient Health Questionnaire 2 (PHQ-2) in identifying major depression in older people. *Journal of the American Geriatrics Society*. 2007;55:596-602.
- (351) Licht-Strunk E, van der Kooij KG, van Schaik DJF et al. Prevalence of depression in older patients consulting their general practitioner in The Netherlands. *International Journal of Geriatric Psychiatry*. 2005;20:1013-1019.
- (352) Licht-Strunk E, Beekman ATF, de Haan M, van Marwijk HWJ. The prognosis of undetected depression in older general practice patients. A one year follow-up study. *Journal of Affective Disorders*. 2009;114:310-315.
- (353) Lichtman JH, Bigger JT, Jr., Blumenthal JA et al. Depression and Coronary Heart Disease. Recommendations for Screening, Referral, and Treatment. A Science Advisory From the American Heart Association Prevention Committee of the Council on Cardiovascular Nursing, Council on Clinical Cardiology, Council on Epidemiology and Prevention, and Interdisciplinary Council on Quality of Care and Outcomes Research. *Circulation*. 2008.
- (354) Lin EHB, Heckbert SR, Rutter CM et al. Depression and Increased Mortality in Diabetes: Unexpected Causes of Death. *Annals of Family Medicine*. 2009;7:414-421.
- (355) Linzer M, Spitzer R, Kroenke K et al. Gender, quality of life, and mental disorders in primary care: Results from the PRIME-MD 1000 study. *Amer J Med*. 1996;101:526-533.
- (356) Liu CF, Bolkan C, Chan D, Yano EM, Rubenstein LV, Chaney EF. Dual Use of VA and Non-VA Services Among Primary Care Patients with Depression. *Journal of General Internal Medicine*. 2009;24:305-311.
- (357) Loerch B, Szegedi A, Kohnen R, Benkert O. The Primary Care Evaluation of Mental Disorders (PRIME-MD), German version: a comparison with the CIDI. *J Psychiat Res*. 2000;34:211-220.
- (358) Loh A, Leonhart R, Wills CE, Simon D, Harter M. The impact of patient participation on adherence and clinical outcome in primary care of depression. *Patient Education and Counseling*. 2007;65:69-78.
- (359) Loh A, Simon D, Wills CE, Kriston L, Niebling W, Harter M. The effects of a shared decision-making intervention in primary care of depression: A cluster-randomized controlled trial. *Patient Education and Counseling*. 2007;67:324-332.

- (360) Lossnitzer N, Muller-Tasch T, Lowe B et al. Exploring Potential Associations of Suicidal Ideation and Ideas of Self-Harm in Patients with Congestive Heart Failure. *Depression and Anxiety*. 2009;26:764-768.
- (361) Lotrakul M, Sumrithe S, Saipanish R. Reliability and validity of the Thai version of the PHQ-9. *Bmc Psychiatry*. 2008;8.
- (362) Lovell K, Bower P, Richards D et al. Developing guided self-help for depression using the Medical Research Council complex interventions framework: a description of the modelling phase and results of an exploratory randomised controlled trial. *Bmc Psychiatry*. 2008;8.
- (363) Lowe B, Grafe K, Kroenke K et al. Predictors of psychiatric comorbidity in medical outpatients. *Psychosomatic Medicine*. 2003;65:764-770.
- (364) Lowe B, Grafe K, Zipfel S et al. Detecting panic disorder in medical and psychosomatic outpatients - Comparative validation of the Hospital Anxiety and Depression Scale, the Patient Health Questionnaire, a screening question, and physicians' diagnosis. *Journal of Psychosomatic Research*. 2003;55:515-519.
- (365) Lowe B, Grafe K, Ufer C et al. Anxiety and depression in patients with pulmonary hypertension. *Psychosomatic Medicine*. 2004;66:831-836.
- (366) Lowe B, Unutzer J, Callahan CM, Perkins AJ, Kroenke K. Monitoring depression treatment outcomes with the patient health questionnaire-9. *Medical Care*. 2004;42:1194-1201.
- (367) Lowe B, Willand L, Eich W et al. Psychiatric comorbidity and work disability in patients with inflammatory rheumatic diseases. *Psychosomatic Medicine*. 2004;66:395-402.
- (368) Lowe B, Spitzer RL, Grafe K et al. Comparative validity of three screening questionnaires for DSM-IV depressive disorders and physicians' diagnoses. *Journal of Affective Disorders*. 2004;78:131-140.
- (369) Lowe B, Grafe K, Zipfel S, Witte S, Loerch B, Herzog W. Diagnosing ICD-10 depressive episodes: Superior criterion validity of the patient health questionnaire. *Psychotherapy and Psychosomatics*. 2004;73:386-390.
- (370) Lowe B, Kroenke K, Herzog W, Grafe K. Measuring depression outcome with a brief self-report instrument: sensitivity to change of the Patient Health Questionnaire (PHQ-9). *Journal of Affective Disorders*. 2004;81:61-66.
- (371) Lowe B, Schenkel I, Bair MJ, Gobel C. Efficacy, predictors of therapy response, and safety of sertraline in routine clinical practice: Prospective, open-label, non-interventional postmarketing surveillance study in 1878 patients. *Journal of Affective Disorders*. 2005;87:271-279.
- (372) Lowe B, Kroenke K, Grafe K. Detecting and monitoring depression with a two-item questionnaire (PHQ-2). *Journal of Psychosomatic Research*. 2005;58:163-171.
- (373) Lowe B, Schenkel I, Carney-Doebbeling C, Gobel C. Responsiveness of the PHQ-9 to psychopharmacological depression treatment. *Psychosomatics*. 2006;47:62-67.

- (374) Lowe B, Decker O, Muller S et al. Validation and standardization of the generalized anxiety disorder screener (GAD-7) in the general population. *Medical Care*. 2008;46:266-274.
- (375) Lowe B, Spitzer RL, Williams JBW, Mussell M, Schellberg D, Kroenke K. Depression, anxiety and somatization in primary care: syndrome overlap and functional impairment. *General Hospital Psychiatry*. 2008;30:191-199.
- (376) Lubetkin EI, Jia HM, Gold MR. Depression, anxiety, and associated health status in low-income Chinese patients. *American Journal of Preventive Medicine*. 2003;24:354-360.
- (377) Luciano JV, yuso-Mateos JL, Fernandez A et al. Utility of the twelve-item World Health Organization Disability Assessment Schedule II (WHO-DAS II) for discriminating depression "caseness" and severity in Spanish primary care patients. *Quality of Life Research*. 2010;19:97-101.
- (378) Luciano JV, yuso-Mateos JL, Fernandez A, Serrano-Blanco A, Roca M, Haro JM. Psychometric properties of the twelve item World Health Organization Disability Assessment Schedule II (WHO-DAS II) in Spanish primary care patients with a first major depressive episode. *Journal of Affective Disorders*. 2010;121:52-58.
- (379) Ludman EJ, Katon W, Russo J et al. Depression and diabetes symptom burden. *General Hospital Psychiatry*. 2004;26:430-436.
- (380) Ludman EJ, Russo JE, Katon WJ et al. How Does Change in Depressive Symptomatology Influence Weight Change in Patients With Diabetes? Observational Results From the Pathways Longitudinal Cohort. *Journals of Gerontology Series A-Biological Sciences and Medical Sciences*. 2010;65:93-98.
- (381) Lynch DJ, McGrady A, Alvarez E, Forman J. Recent life changes and medical utilization in an academic family practice. *Journal of Nervous and Mental Disease*. 2005;193:633-635.
- (382) Macran S, Kind P, Collingwood J, Hull R, McDonald I, Parkinson L. Evaluating podiatry services: Testing a treatment specific measure of health status. *Quality of Life Research*. 2003;12:177-188.
- (383) Maguen S, Lucenko BA, Reger MA et al. The Impact of Reported Direct and Indirect Killing on Mental Health Symptoms in Iraq War Veterans. *Journal of Traumatic Stress*. 2010;23:86-90.
- (384) Maizels M, Burchette R. Somatic symptoms in headache patients: The influence of headache diagnosis, frequency, and comorbidity. *Headache*. 2004;44:983-993.
- (385) Maizels M, Smitherman TA, Penzien DB. A review of screening tools for psychiatric comorbidity in headache patients. *Headache*. 2006;46:S98-S109.
- (386) Maizels M, Houle T. Results of screening with the brief headache screen compared with a modified ID Migraine (TM). *Headache*. 2008;48:385-394.
- (387) Mallik S, Spertus JA, Reid KJ et al. Depressive symptoms after acute myocardial infarction - Evidence for highest rates in younger women. *Archives of Internal Medicine*. 2006;166:876-883.

- (388) Malmenstrom M, Bixo M, Bjorn I, Astrom M, Poromaa IS. Patients with psychiatric disorders in gynecologic practice - a three year follow-up. *Journal of Psychosomatic Obstetrics and Gynecology*. 2006;27:17-22.
- (389) Mann R, Gilbody S, Richards D. Putting the 'Q' in depression QALYs: a comparison of utility measurement using EQ-5D and SF-6D health related quality of life measures. *Social Psychiatry and Psychiatric Epidemiology*. 2009;44:569-578.
- (390) Mao LM, Kidd MR, Rogers G et al. Social factors associated with Major Depressive Disorder in homosexually active, gay men attending general practices in urban Australia. *Australian and New Zealand Journal of Public Health*. 2009;33:83-86.
- (391) Marchesi C, Bertoni S, Cantoni A, Maggini C. Is alexithymia a personality trait increasing the risk of depression? A prospective study evaluating alexithymia before, during and after a depressive episode. *Psychological Medicine*. 2008;38:1717-1722.
- (392) Margaretten M, Yelin E, Imboden J et al. Predictors of Depression in a Multiethnic Cohort of Patients With Rheumatoid Arthritis. *Arthritis & Rheumatism-Arthritis Care & Research*. 2009;61:1586-1591.
- (393) Mario B, Massimiliano M, Chiara M, Alessandro S, Antonella C, Gianfranco F. White-coat effect among older patients with suspected cognitive impairment: prevalence and clinical implications. *International Journal of Geriatric Psychiatry*. 2009;24:509-517.
- (394) Marlow RA, Kegowicz CL, Starkey KN. Prevalence of Depression Symptoms in Outpatients with a Complaint of Headache. *Journal of the American Board of Family Medicine*. 2009;22:633-637.
- (395) Marsh J, Patel S, Gelaye B et al. Prevalence of Workplace Abuse and Sexual Harassment among Female Faculty and Staff. *Journal of Occupational Health*. 2009;51:314-322.
- (396) Martin A, Rief W, Klaiberg A, Braehler E. Validity of the Brief Patient Health Questionnaire Mood Scale (PHQ-9) in the general population. *General Hospital Psychiatry*. 2006;28:71-77.
- (397) Mata S, Gonzalez A, Lavie R, Resler G. Validation of the PRIME-MD for the detection of generalized anxiety disorder. *Investigacion Clinica*. 2008;49:39-48.
- (398) Mauksch LB, Tucker SM, Katon WJ et al. Mental illness, functional impairment, and patient preferences for collaborative care in an uninsured, primary care population. *J Fam Pract*. 2001;50:41-47.
- (399) Mauksch LB, Katon WJ, Russo J, Tucker SM, Walker E, Cameron J. The content of a low-income, uninsured primary care population: Including the patient agenda. *Journal of the American Board of Family Practice*. 2003;16:278-289.
- (400) McCarrier KP, Ralston JD, Hirsch IB et al. Web-Based Collaborative Care for Type 1 Diabetes: A Pilot Randomized Trial. *Diabetes Technology & Therapeutics*. 2009;11:211-217.
- (401) McClave AK, Dube SR, Strine TW, Kroenke K, Caraballo RS, Mokdad AH. Associations between smoking cessation and anxiety and depression among US adults. *Addictive Behaviors*. 2009;34:491-497.

- (402) McGuire LC, Strine TW, Vachirasudlekha S, Mokdad AH, Anderson LA. The prevalence of depression in older US women: 2006 Behavioral Risk Factor Surveillance System. *Journal of Womens Health*. 2008;17:501-507.
- (403) McGuire LC, Strine TW, Allen RS, Anderson LA, Mokdad AH. The Patient Health Questionnaire 8: Current Depressive Symptoms Among US Older Adults, 2006 Behavioral Risk Factor Surveillance System. *American Journal of Geriatric Psychiatry*. 2009;17:324-334.
- (404) McLaughlin TJ, Aupont O, Bambauer KZ et al. Improving psychologic adjustment to chronic illness in cardiac patients - The role of depression and anxiety. *Journal of General Internal Medicine*. 2005;20:1084-1090.
- (405) McManus D, Pipkin SS, Whooley MA. Screening for depression in patients with coronary heart disease (data from the heart and soul study). *American Journal of Cardiology*. 2005;96:1076-1081.
- (406) McNutt LA, Carlson BE, Persaud M, Postmus J. Cumulative abuse experiences, physical health and health behaviors. *Ann Epidemiol*. 2002;12:123-130.
- (407) McRae AL, Brady KT, Sonne SC. Alcohol and substance abuse. *Med Clin N Amer*. 2001;85:779-+.
- (408) Means-Christensen AJ, Arnau RC, Tonidandel AM, Bramson R, Meagher MW. An efficient method of identifying major depression and panic disorder in primary care. *Journal of Behavioral Medicine*. 2005;28:565-572.
- (409) Melville JL, Miller EA, Fialkow MF, Lentz GM, Miller JL, Fenner DE. Relationship between patient report and physician assessment of urinary incontinence severity. *American Journal of Obstetrics and Gynecology*. 2003;189:76-80.
- (410) Meshberg-Cohen S, Svikis D. Panic disorder, trait anxiety, and alcohol use in pregnant and nonpregnant women. *Comprehensive Psychiatry*. 2007;48:504-510.
- (411) Mewes R, Rief W, Brahler E, Martin A, Glaesmer H. Lower decision threshold for doctor visits as a predictor of health care use in somatoform disorders and in the general population. *General Hospital Psychiatry*. 2008;30:349-355.
- (412) Mewes R, Rief W, Stenzel N, Glaesmer H, Martin A, Brahler E. What is "normal" disability? An investigation of disability in the general population. *Pain*. 2009;142:36-41.
- (413) Mikolajczyk E, Zietek J, Samochowiec A, Samochowiec J. Personality dimensions measured using the Temperament and Character Inventory (TCI) and NEO-FFI on a Polish sample. *International Journal of Methods in Psychiatric Research*. 2008;17:210-219.
- (414) Millette K, Hudson M, Baron M, Thombs BD. Comparison of the PHQ-9 and CES-D depression scales in systemic sclerosis: internal consistency reliability, convergent validity and clinical correlates. *Rheumatology*. 2010;49:789-796.
- (415) Miranda J, Azocar F, Komaromy M, Golding JM. Unmet mental health needs of women in public-sector gynecologic clinics. *Am J Obstet Gynecol*. 1998;178:212-217.

- (416) Miranda J, Siddique J, Belin TR, Kohn-Wood LP. Depression prevalence in disadvantaged young black women - African and Caribbean immigrants compared to US-born African Americans. *Social Psychiatry and Psychiatric Epidemiology*. 2005;40:253-258.
- (417) Mitchell AJ, McGlinchey JB, Young D, Chelminski I, Zimmerman M. Accuracy of specific symptoms in the diagnosis of major depressive disorder in psychiatric out-patients: data from the MIDAS project. *Psychological Medicine*. 2009;39:1107-1116.
- (418) Mohr DC, Hart SL, Howard I et al. Barriers to psychotherapy among depressed and nondepressed primary care patients. *Annals of Behavioral Medicine*. 2006;32:254-258.
- (419) Monahan PO, Kroenke K. PHQ-9 and PHQ-2 in Western Kenya. *Journal of General Internal Medicine*. 2009;24:891.
- (420) Monahan PO, Shacham E, Reece M et al. Validity/reliability of PHQ-9 and PHQ-2 depression scales among adults living with HIV/AIDS in western Kenya. *J Gen Intern Med*. 2009;24:189-197.
- (421) Muhtz C, Zyriax BC, Klahn T, Windler E, Otte C. Depressive symptoms and metabolic risk: Effects of cortisol and gender. *Psychoneuroendocrinology*. 2009;34:1004-1011.
- (422) Muller-Tasch T, Peters-Klimm F, Schellberg D et al. Depression is a major determinant of quality of life in patients with chronic systolic heart failure in general practice. *Journal of Cardiac Failure*. 2007;13:818-824.
- (423) Mulrow CD, Williams JW, Jr., Gerety MB, Ramirez G, Montiel OM, Kerber C. Case-finding instruments for depression in primary care settings. *Ann Intern Med*. 1995;122:913-921.
- (424) Muramatsu K, Miyaoka H, Kamijima K et al. The patient health questionnaire, Japanese version: Validity according to the mini-international neuropsychiatric interview-plus. *Psychological Reports*. 2007;101:952-960.
- (425) Mussell M, Kroenke K, Spitzer RL, Williams JBW, Herzog W, Lowe B. Gastrointestinal symptoms in primary care: Prevalence and association with depression and anxiety. *Journal of Psychosomatic Research*. 2008;64:605-612.
- (426) Na YM, Kim KS, Lee KU et al. The relationship between depressive symptoms in outpatients with chronic illness and health care costs. *Yonsei Medical Journal*. 2007;48:787-794.
- (427) Nau DP, Aikens JE, Pacholski AM. Effects of gender and depression on oral medication adherence in persons with type 2 diabetes mellitus. *Gender Medicine*. 2007;4:205-213.
- (428) Nazemi H, Larkin AA, Sullivan MD, Katon W. Methodological issues in the recruitment of primary care patients with depression. *Int J Psychiat Med*. 2001;31:277-288.
- (429) Nease DE, Klinkman MS, Volk RJ. Improved detection of depression in primary care through severity evaluation. *Journal of Family Practice*. 2002;51:1065-1070.
- (430) Nease DE, Klinkman MS, Aikens JE. Severity and criteria based prompting for treatment of depressed patients. *International Journal of Psychiatry in Medicine*. 2005;35:149-159.

- (431) Nease DE, Jr., Volk RJ, Cass AR. Does the severity of mood and anxiety symptoms predict health care utilization? *J Fam Pract.* 1999;48:769-777.
- (432) Nease DE, Jr., Volk RJ, Cass AR. Investigation of a severity-based classification of mood and anxiety symptoms in primary care patients. *J Am Board Fam Pract.* 1999;12:21-31.
- (433) Nelson DB, Murdoch M, Sandozi IK, Dalmaso AP, Crabtree JE, Ho SB. Dyspepsia is associated with CagA-positive *Helicobacter pylori*. *Amer J Gastroenterol.* 2000;95:3412-3417.
- (434) Neria Y, Gross R, Olfson M et al. Posttraumatic stress disorder in primary care one year after the 9/11 attacks. *General Hospital Psychiatry.* 2006;28:213-222.
- (435) Neria Y, Olfson M, Gameroff MJ et al. The Mental Health Consequences of Disaster-Related Loss: Findings from Primary Care One Year After the 9/11 Terrorist Attacks. *Psychiatry-Interpersonal and Biological Processes.* 2008;71:339-348.
- (436) Norton J, De Roquefeuil G, Boulenger JP, Ritchie K, Mann A, Tylee A. Use of the PRIME-MD Patient Health Questionnaire for estimating the prevalence of psychiatric disorders in French primary care: comparison with family practitioner estimates and relationship to psychotropic medication use. *General Hospital Psychiatry.* 2007;29:285-293.
- (437) Norton J, De Roquefeuil G, David M, Boulenger JP, Ritchie K, Mann A. Prevalence of psychiatric disorders in French general practice using the patient health questionnaire: Comparison with GP case-recognition and psychotropic medication prescription. *Encephale-Revue de Psychiatrie Clinique Biologique et Therapeutique.* 2009;35:560-569.
- (438) O'Malley PG, Jackson JL, Kroenke K, Yoon IK, Hornstein E, Dennis GJ. The value of screening for psychiatric disorders in rheumatology referrals. *Arch Intern Med.* 1998;158:2357-2362.
- (439) O'Malley PG, Wong PWK, Kroenke K, Roy MJ, Wong RKH. The value of screening for psychiatric disorders prior to upper endoscopy. *J Psychosom Res.* 1998;44:279-287.
- (440) Odden MC, Whooley MA, Shlipak MG. Depression, stress, and quality of life in persons with chronic kidney disease: The Heart and Soul Study. *Nephron Clinical Practice.* 2006;103:C1-C7.
- (441) Okulate G, Olayinka O, Dogunro AS. Erectile dysfunction: prevalence and relationship to depression, alcohol abuse and panic disorder. *General Hospital Psychiatry.* 2003;25:209-213.
- (442) Okulate GT, Olayinka MO, Jones OBE. Somatic symptoms in depression: evaluation of their diagnostic weight in an African setting. *British Journal of Psychiatry.* 2004;184:422-427.
- (443) Olfson M, Shea S, Feder A et al. Prevalence of anxiety, depression, and substance use disorders in an urban general medicine practice. *Arch Fam Med.* 2000;9:876-883.
- (444) Olfson M, Gameroff MJ. Generalized anxiety disorder, somatic pain and health care costs. *General Hospital Psychiatry.* 2007;29:310-316.
- (445) Olson AL, Dietrich AJ, Prazar G, Hurley J. Brief maternal depression screening at well-child visits. *Pediatrics.* 2006;118:207-216.

- (446) Omoro SAO, Fann JR, Weymuller EA, Macharia IM, Yueh B. Swahili translation and validation of the patient health questionnaire-9 depression scale in the Kenyan head and neck cancer patient population. *International Journal of Psychiatry in Medicine*. 2006;36:367-381.
- (447) Osborne TL, Turner AP, Williams RM et al. Correlates of pain interference in multiple sclerosis. *Rehabilitation Psychology*. 2006;51:166-174.
- (448) Osorio FD, Mendes AV, Crippa JA, Loureiro SR. Study of the Discriminative Validity of the PHQ-9 and PHQ-2 in a Sample of Brazilian Women in the Context of Primary Health Care. *Perspectives in Psychiatric Care*. 2009;45:216-227.
- (449) Otte C, Marmar CR, Pipkin SS, Moos R, Browner WS, Whooley MA. Depression and 24-hour urinary cortisol in medical outpatients with coronary heart disease: The Heart and Soul Study. *Biol Psychiatry*. 2004;56:241-247.
- (450) Otte C, Neylan TC, Pipkin SS, Browner WS, Whooley MA. Depressive symptoms and 24-hour urinary norepinephrine excretion levels in patients with coronary disease: Findings from the heart and soul study. *American Journal of Psychiatry*. 2005;162:2139-2145.
- (451) Pang AHT, Chao DVK, Fabb WE, Lai KYC, Leung T. Validation study of the chinese version of the Primary Care Evaluation of Mental Disorders (cPRIME-MD). Part II - Validity. *MediMedia Asia*. 1997.
- (452) Parashar S, Rumsfeld JS, Spertus JA et al. Time course of depression and outcome of myocardial infarction. *Archives of Internal Medicine*. 2006;166:2035-2043.
- (453) Parker T, May PA, Maviglia MA, Petrakis S, Sunde S, Gloyd SV. PRIME-MD: Its utility in detecting mental disorders in American Indians. *Int J Psychiat Med*. 1997;27:107-128.
- (454) Patel V, Araya R, Chowdhary N et al. Detecting common mental disorders in primary care in India: a comparison of five screening questionnaires. *Psychological Medicine*. 2008;38:221-228.
- (455) Patten SB, Schopflocher D. Longitudinal epidemiology of major depression as assessed by the Brief Patient Health Questionnaire (PHQ-9). *Comprehensive Psychiatry*. 2009;50:26-33.
- (456) Pavlik VN, Hyman DJ, Wendt JA, Orengo C. Association of a culturally defined syndrome (Nervios) with chest pain and DSM-IV affective disorders in Hispanic patients referred for cardiac stress testing. *Ethnicity & Disease*. 2004;14:505-514.
- (457) Pedersen SS, Denollet J, De Jonge P, Simsek C, Serruys PW, van Domburg RT. Brief Depression Screening with the PHQ-2 Associated with Prognosis Following Percutaneous Coronary Intervention with Paclitaxel-Eluting Stenting. *Journal of General Internal Medicine*. 2009;24:1037-1042.
- (458) Pedrazzini B, Waldvogel S, Cornuz J et al. The impact of iron supplementation efficiency in female blood donors with a decreased ferritin level and no anaemia. Rationale and design of a randomised controlled trial: a study protocol. *Trials*. 2009;10.
- (459) Peele PB, Tollerud DJ. Depression and occupational injury: Results of a pilot investigation. *Journal of Occupational and Environmental Medicine*. 2005;47:424-427.

- (460) Persoons P, Luyckx K, Desloovere C, Vandenberghe J, Fischler B. Anxiety and mood disorders in otorhinolaryngology outpatients presenting with dizziness: Validation of the self-administered PRIME-MD Patient Health Questionnaire and epidemiology. *General Hospital Psychiatry*. 2003;25:316-323.
- (461) Persoons P, Vermeire S, Demyttenaere K et al. The impact of major depressive disorder on the short- and long-term outcome of Crohn's disease treatment with infliximab. *Alimentary Pharmacology & Therapeutics*. 2005;22:101-110.
- (462) Peterlin BL, Tietjen GE, Brandes JL et al. Posttraumatic Stress Disorder in Migraine. *Headache*. 2009;49:541-551.
- (463) Peters-Klimm F, Campbell S, Muller-Tasch T et al. Primary care-based multifaceted, interdisciplinary medical educational intervention for patients with systolic heart failure: lessons learned from a cluster randomised controlled trial. *Trials*. 2009;10.
- (464) Philbrick JT, Connelly JE, Wofford AB. The prevalence of mental disorders in rural office practice. *J Gen Intern Med*. 1996;11:9-15.
- (465) Picardi A, Amerio P, Baliva G et al. Recognition of depressive and anxiety disorders in dermatological outpatients. *Acta Dermato-Venereologica*. 2004;84:213-217.
- (466) Picardi A, Adler DA, Abeni D et al. Screening for depressive disorders in patients with skin diseases: A comparison of three screeners. *Acta Dermato-Venereologica*. 2005;85:414-419.
- (467) Picardi A, Mazzotti E, Pasquini P. Prevalence and correlates of suicidal ideation among patients with skin disease. *Journal of the American Academy of Dermatology*. 2006;54:420-426.
- (468) Pilowsky DJ, Olfson M, Gameroff MJ et al. Panic disorder and suicidal ideation in primary care. *Depression and Anxiety*. 2006;23:11-16.
- (469) Pinto-Meza A, Serrano-Blanco A, Penarrubia MT, Blanco E, Haro JM. Assessing depression in primary care with the PHQ-9: Can it be carried out over the telephone? *Journal of General Internal Medicine*. 2005;20:738-742.
- (470) Pinto-Meza A, Usall J, Serrano-Blanco A, Suarez D, Haro JM. Gender differences in response to antidepressant treatment prescribed in primary care. Does menopause make a difference? *Journal of Affective Disorders*. 2006;93:53-60.
- (471) Pinto-Meza A, Fernandez A, Serrano-Blanco A, Haro JM. Adequacy of antidepressant treatment in Spanish primary care: A naturalistic six-month follow-up study. *Psychiatric Services*. 2008;59:78-83.
- (472) Pirkis J, Pfaff J, Williamson M et al. The community prevalence of depression in older Australians. *Journal of Affective Disorders*. 2009;115:54-61.
- (473) Poleshuck EL, Bair MJ, Kroenke K, Watts A, Tu X, Giles DE. Pain and Depression in Gynecology Patients. *Psychosomatics*. 2009;50:270-276.

- (474) Poongothai S, Pradeepa R, Ganesan A, Mohan V. Prevalence of Depression in a Large Urban South Indian Population - The Chennai Urban Rural Epidemiology Study (Cures-70). *Plos One*. 2009;4.
- (475) Porcerelli JH, Huprich SK, Binienda J, Karana D. Object representations and their relationship to psychopathology and physical health status in African-American women in primary care. *Journal of Nervous and Mental Disease*. 2006;194:838-844.
- (476) Porcerelli JH, Olson TR, Presniak MD, Markova T, Miller K. Defense Mechanisms and Major Depressive Disorder in African American Women. *Journal of Nervous and Mental Disease*. 2009;197:736-741.
- (477) Porcerelli JH, Huprich SK, Markova T. Mental Representations in Women With Panic Disorder An Urban African-American Sample. *Journal of Nervous and Mental Disease*. 2010;198:144-149.
- (478) Pratt LA, Brody DJ. Depression in the United States household population, 2005-2006. *NCHS Data Brief*. 2008;7.
- (479) Preville M, Cote G, Boyer R, Hebert R. Detection of depression and anxiety disorders by home care nurses. *Aging & Mental Health*. 2004;8:400-409.
- (480) Price SK, Proctor EK. A Rural Perspective on Perinatal Depression: Prevalence, Correlates, and Implications for Help-Seeking Among Low-income Women. *Journal of Rural Health*. 2009;25:158-166.
- (481) Pyne JM, Fortney JC, Tripathi S, Feeny D, Ubel P, Brazier J. How Bad Is Depression? Preference Score Estimates from Depressed Patients and the General Population. *Health Services Research*. 2009;44:1406-1423.
- (482) Ramirez MTG, Hernandez RL, Garcia-Campayo J. The relationship among depression, anxiety, and somatic symptoms in a sample of university students in northern Mexico. *Revista Panamericana de Salud Publica-Pan American Journal of Public Health*. 2009;25:141-145.
- (483) Rasmussen NH, Bernard ME, Harmsen WS. Physical symptoms that predict psychiatric disorders in rural primary care adults. *Journal of Evaluation in Clinical Practice*. 2008;14:399-406.
- (484) Rasmussen S. "Prime-MD"--a rapid method for diagnosis of mental disease in general practice. *Ugeskr Laeger*. 1998;160:5672.
- (485) Reed SD, Ludman EJ, Newton KM et al. Depressive symptoms and menopausal burden in the midlife. *Maturitas*. 2009;62:306-310.
- (486) Reger MA, Gahm GA, Swanson RD, Duma SJ. Association Between Number of Deployments to Iraq and Mental Health Screening Outcomes in US Army Soldiers. *Journal of Clinical Psychiatry*. 2009;70:1266-1272.
- (487) Rentsch D, Dumont P, Borgacci S et al. Prevalence and treatment of depression in a hospital department of internal medicine. *General Hospital Psychiatry*. 2007;29:25-31.

- (488) Reuber M, Burness C, Howlett S, Brazier J, Grunewald R. Tailored psychotherapy for patients with functional neurological symptoms: A pilot study. *Journal of Psychosomatic Research*. 2007;63:625-632.
- (489) Reuland DS, Cherrington A, Watkins GS, Bradford DW, Blanco RA, Gaynes BN. Diagnostic Accuracy of Spanish Language Depression-Screening Instruments. *Annals of Family Medicine*. 2009;7:455-462.
- (490) Reuter M, Jecht M, Bierwirth RA et al. Diabetic Polyneuropathy and Related Comorbidities: Results of the GloDi Project. *Diabetologie und Stoffwechsel*. 2009;4:230-237.
- (491) Revicki DA, Brandenburg N, Matza L, Hornbrook MC, Feeny D. Health-related quality of life and utilities in primary-care patients with generalized anxiety disorder. *Quality of Life Research*. 2008;17:1285-1294.
- (492) Rhee MK, Musselman D, Ziemer DC et al. Unrecognized glucose intolerance is not associated with depression. Screening for Impaired Glucose Tolerance study 3 (SIGT 3). *Diabetic Medicine*. 2008;25:1361-1365.
- (493) Richards DA, Lovell K, Gilbody S et al. Collaborative care for depression in UK primary care: a randomized controlled trial. *Psychological Medicine*. 2008;38:279-287.
- (494) Richards DA, Suckling R. Improving access to psychological therapies: Phase IV prospective cohort study. *British Journal of Clinical Psychology*. 2009;48:377-396.
- (495) Richards DA, Hughes-Morley A, Hayes RA et al. Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depression - study protocol. *Bmc Health Services Research*. 2009;9.
- (496) Richardson EJ, Richards JS. Factor structure of the PHQ-9 screen for depression across time since injury among persons with spinal cord injury. *Rehabilitation Psychology*. 2008;53:243-249.
- (497) Richardson L, McCauley E, Katon W. Collaborative care for adolescent depression: a pilot study. *General Hospital Psychiatry*. 2009;31:36-45.
- (498) Richardson LP, Katzenellengoben R. Childhood and adolescent depression: The role of primary care providers in diagnosis and treatment. *Current Problems in Pediatric and Adolescent Health Care*. 2005;35:6-24.
- (499) Riddle DL, Wade JB, Jiranek WA, Kong XR. Preoperative Pain Catastrophizing Predicts Pain Outcome after Knee Arthroplasty. *Clinical Orthopaedics and Related Research*. 2010;468:798-806.
- (500) Riddle JR, Smith TC, Smith B et al. Millennium Cohort: the 2001-2003 baseline prevalence of mental disorders in the U.S. military. *J Clin Epidemiol*. 2007;60:192-201.
- (501) Rief W, Nanke A, Klaiberg A, Braehler E. Base rates for panic and depression according to the Brief Patient Health Questionnaire: a population-based study. *J Affect Disord*. 2004;82:271-276.

- (502) Rief W, Martin A, Klaiberg A, Brahler E. Specific effects of depression, panic, and somatic symptoms on illness behavior. *Psychosomatic Medicine*. 2005;67:596-601.
- (503) Robison J, Gruman C, Gaztambide S, Blank K. Screening for depression in middle-aged and older Puerto Rican primary care patients. *J Gerontol Ser A-Biol Sci Med*. 2002;57:M308-M314.
- (504) Roca M, Gili M, Garcia-Garcia M et al. Prevalence and comorbidity of common mental disorders in primary care. *Journal of Affective Disorders*. 2009;119:52-58.
- (505) Rogal SS, Poschman K, Belanger K et al. Effects of posttraumatic stress disorder on pregnancy outcomes. *Journal of Affective Disorders*. 2007;102:137-143.
- (506) Rogers WH, Wilson IB, Bungay KM, Cynn DJ, Adler DA. Assessing the performance of a new depression screener for primary care (PC-SAD (c)). *J Clin Epidemiol*. 2002;55:164-175.
- (507) Rogers WH, Adler DA, Bungay KM, Wilson IB. Depression screening instruments made good severity measures in a cross-sectional analysis. *Journal of Clinical Epidemiology*. 2005;58:370-377.
- (508) Rollman BL, Hanusa BH, Gilbert T, Lowe HJ, Kapoor WN, Schulberg HC. The electronic medical record. *Arch Intern Med*. 2001;161:189-197.
- (509) Rollman BL, Hanusa BH, Lowe HJ, Gilbert T, Kapoor WN, Schulberg HC. A randomized trial using computerized decision support to improve treatment of major depression in primary care. *J Gen Intern Med*. 2002;17:493-503.
- (510) Rollman BL, Belnap BH, Mazumdar S et al. Symptomatic severity of PRIME-MD diagnosed episodes of panic and generalized anxiety disorder in primary care. *Journal of General Internal Medicine*. 2005;20:623-628.
- (511) Rollman BL, Belnap BH, LeMenager MS, Mazumdar S, Schulberg HC, Reynolds CF. The Bypassing the Blues Treatment Protocol: Stepped Collaborative Care for Treating Post-CABG Depression. *Psychosomatic Medicine*. 2009;71:217-230.
- (512) Rollman BL, Belnap BH, Mazumdar S et al. A Positive Phq-2 Depression Screen Among Hospitalized Chf Patients Predicts Cardiovascular Morbidity at 6-Months Follow-Up. *Journal of General Internal Medicine*. 2009;24:8.
- (513) Rosemann T, Korner T, Wensing M et al. Rationale, design and conduct of a comprehensive evaluation of a primary care based intervention to improve the quality of life of osteoarthritis patients. The PraxArt-project: a cluster randomized controlled trial. *Bmc Public Health*. 2005;5:77.
- (514) Rosemann T, Kuehle T, Laux G, Szecsenyi J. Osteoarthritis of the knee and hip: a comparison of factors associated with physical activity. *Clinical Rheumatology*. 2007;26:1811-1817.
- (515) Rosemann T, Laux G, Kuehle T. Osteoarthritis and functional disability: results of a cross sectional study among primary care patients in Germany. *Bmc Musculoskeletal Disorders*. 2007;8.

- (516) Rosemann T, Joos S, Szecsenyi J, Laux G, Wensing M. Health service utilization patterns of primary care patients with osteoarthritis. *Bmc Health Services Research*. 2007;7.
- (517) Rosemann T, Backenstrass M, Joest K, Rosemann A, Szecsenyi J, Laux G. Predictors of depression in a sample of 1,021 primary care patients with osteoarthritis. *Arthritis & Rheumatism-Arthritis Care & Research*. 2007;57:415-422.
- (518) Rosemann T, Grol R, Herman K, Wensing M, Szecsenyi J. Association between obesity, quality of life, physical activity and health service utilization in primary care patients with osteoarthritis. *International Journal of Behavioral Nutrition and Physical Activity*. 2008;5.
- (519) Rosemann T, Laux G, Szecsenyi J, Grol R. The Chronic Care Model: congruency and predictors among primary care patients with osteoarthritis. *Quality & Safety in Health Care*. 2008;17:442-446.
- (520) Rosemann T, Laux G, Szecsenyi J, Wensing M, Grol R. Pain and osteoarthritis in primary care: Factors associated with pain perception in a sample of 1,021 patients. *Pain Medicine*. 2008;9:903-910.
- (521) Rosemann T, Wensing M, Szecsenyi J, Grol R. Satisfaction of osteoarthritis patients with provided care is not related to the disease-specific quality of life. *Journal of Evaluation in Clinical Practice*. 2009;15:486-491.
- (522) Rubio-Valera M, Serrano-Blanco A, Trave P, Penarrubia-Maria T, Ruiz M, Pujol MM. Community pharmacist intervention in depressed primary care patients (PRODEFAR study): randomized controlled trial protocol. *Bmc Public Health*. 2009;9.
- (523) Ruiz MA, Zamorano E, Garcia-Campayo J, Pardo A, Freire O, Rejas J. Further Validity Evidences of the Generalized Anxiety Disorder-7 Scale (Gad-7). *Value in Health*. 2009;12:A362.
- (524) Rumsfeld JS, Ho PM. Depression and cardiovascular disease: a call for recognition. *Circulation*. 2005;111:250-253.
- (525) Ruo B, Rumsfeld JS, Hlatky MA, Liu HY, Browner WS, Whooley MA. Depressive symptoms and health-related quality of life - The heart and soul study. *Jama-Journal of the American Medical Association*. 2003;290:215-221.
- (526) Ruo B, Rumsfeld JS, Pipkin S, Whooley MA. Relation between depressive symptoms and treadmill exercise capacity in the Heart and Soul Study. *Am J Cardiol*. 2004;94:96-99.
- (527) Ruoff G. A method that dramatically improves patient adherence to depression treatment. *Journal of Family Practice*. 2005;54:846-852.
- (528) Rush AJ, First MB, Blacker D. *Handbook of Psychiatric Measures*. Second Edition ed. Arlington, VA: American Psychiatric Publishing, Inc.; 2007.
- (529) Russell AS, Hui BK. The use of PRIME-MD questionnaire in a rheumatology clinic. *Rheumatol Int*. 2005;25:292-295.

- (530) Ryan MAK, Smith TC, Smith B et al. Millennium Cohort: enrollment begins a 21-year contribution to understanding the impact of military service. *Journal of Clinical Epidemiology*. 2007;60:181-191.
- (531) Sachdev PS, Parslow RA, Lux O et al. Relationship of homocysteine, folic acid and vitamin B-12 with depression in a middle-aged community sample. *Psychological Medicine*. 2005;35:529-538.
- (532) Sakakibara BM, Miller WC, Orenczuk SG, Wolfe DL. A systematic review of depression and anxiety measures used with individuals with spinal cord injury. *Spinal Cord*. 2009;47:841-851.
- (533) Sanders LB. Assessing and managing women with depression: A midwifery perspective. *Journal of Midwifery & Womens Health*. 2006;51:185-192.
- (534) Schenker Y, Stewart A, Na B, Whooley MA. Depressive Symptoms and Perceived Doctor-Patient Communication in the Heart and Soul Study. *Journal of General Internal Medicine*. 2009;24:550-556.
- (535) Schenker Y, Stewart A, Na B, Whooley MA. Depressive Symptoms and Perceived Doctor-Patient Communication in the Heart and Soul Study. *J Gen Intern Med*. 2009.
- (536) Schmidt VM. "Prime-MD"--a quick method for diagnosis of mental disease in general practice. Is the documentation correct? *Ugeskr Laeger*. 1998;160:5370-5371.
- (537) Schmitz N, Kruse J, Tress W. Improving screening for mental disorders in the primary care setting by combining the GHQ-12 and SCL-90-R subscales. *Compr Psychiat*. 2001;42:166-173.
- (538) Schneider A, Biessecker K, Quinzler R et al. Asthma patients with low perceived burden of illness: a challenge for guideline adherence. *Journal of Evaluation in Clinical Practice*. 2007;13:846-852.
- (539) Schneider A, Lowe B, Meyer FJ, Biessecker K, Joos S, Szecsenyi J. Depression and panic disorder as predictors of health outcomes for patients with asthma in primary care. *Respiratory Medicine*. 2008;102:359-366.
- (540) Scholle SH, Haskett RF, Hanusa BH, Pincus HA, Kupfer DJ. Addressing depression in obstetrics/gynecology practice. *General Hospital Psychiatry*. 2003;25:83-90.
- (541) Schomerus G, Matschinger H, Angermeyer MC. Attitudes that determine willingness to seek psychiatric help for depression: a representative population survey applying the Theory of Planned Behaviour. *Psychological Medicine*. 2009;39:1855-1865.
- (542) Schriger DL, Gibbons PS, Langone CA, Lee S, Altshuler LL. Enabling the diagnosis of occult psychiatric illness in the emergency department: A randomized, controlled trial of the computerized, self-administered PRIME-MD diagnostic system. *Ann Emerg Med*. 2001;37:132-140.
- (543) Schroer S, MacPherson H. Acupuncture, or non-directive counselling versus usual care for the treatment of depression: a pilot study. *Trials*. 2009;10.

- (544) Schubert CC, Boustani M, Callahan CM, Perkins AJ, Hui S, Hendrie HC. Acute Care Utilization by Dementia Caregivers Within Urban Primary Care Practices. *Journal of General Internal Medicine*. 2008;23:1736-1740.
- (545) Schulberg HC, Lee PW, Bruce ML et al. Suicidal ideation and risk levels among primary care patients with uncomplicated depression. *Annals of Family Medicine*. 2005;3:523-528.
- (546) Schwenk TL, Gorenflo DW, Leja LM. A survey on the impact of being depressed on the professional status and mental health care of physicians. *Journal of Clinical Psychiatry*. 2008;69:617-620.
- (547) Sears SF, Danda CE, Evans GD. PRIME-MD and rural primary care: Detecting depression in a low- income rural population. *Prof Psychol-Res Pract*. 1999;30:357-360.
- (548) Seminario NA, Farias ST, Jorgensen J, Bourgeois JA, Seyal M. Determination of prevalence of depression in an epilepsy clinic using a brief DSM-IV-based self-report questionnaire. *Epilepsy & Behavior*. 2009;15:362-366.
- (549) Serrano-Blanco A, Pinto-Meza A, Penarrubia MT et al. Selective serotonin reuptake inhibitors: are there differences in effectiveness in primary care? *Primary Care & Community Psychiatry*. 2006;11:113-120.
- (550) Serrano-Blanco A, Pinto-Meza A, Suarez D, Penarrubia MT, Haro JM. Cost-utility of selective serotonin reuptake inhibitors for depression in primary care in Catalonia. *Acta Psychiatrica Scandinavica*. 2006;114:39-47.
- (551) Serrano-Blanco A, Palao DJ, Luciano JV et al. Prevalence of mental disorders in primary care: results from the diagnosis and treatment of mental disorders in primary care study (DASMAP). *Social Psychiatry and Psychiatric Epidemiology*. 2010;45:201-210.
- (552) Sha MC, Callahan CM, Counsell SR, Westmoreland GR, Stump TE, Kroenke K. Physical symptoms as a predictor of health care use and mortality among older adults. *American Journal of Medicine*. 2005;118:301-306.
- (553) Shacham E, Nurutdinova D, Satyanarayana V, Stamm K, Overton ET. Routine Screening for Depression: Identifying a Challenge for Successful HIV Care. *Aids Patient Care and Stds*. 2009;23:949-955.
- (554) Shah MN, Karuza J, Rueckmann E, Swanson P, Conwell Y, Katz P. Reliability and Validity of Prehospital Case Finding for Depression and Cognitive Impairment. *Journal of the American Geriatrics Society*. 2009;57:697-702.
- (555) Sharma V, Zehtabchi S, Rojas N, Birkhahn R. Ethnic Variations in Quality of Life and Depressive Symptoms Among Black Americans With Acute Decompensated Heart Failure. *Journal of the National Medical Association*. 2009;101:985-991.
- (556) Shear K, Belnap BH, Mazumdar S, Houck P, Rollman BL. Generalized anxiety disorder severity scale (GADSS): a preliminary validation study. *Depress Anxiety*. 2006;23:77-82.

- (557) Sheeran T, Reilly C, Pomerantz J, Mlodzianowski A, Bruce M. A Training Program to Screen for Geriatric Depression Using the Phq-9. *Gerontologist*. 2008;48:551.
- (558) Shtayermman O. Peer victimization in adolescents and young adults diagnosed with Asperger's Syndrome: a link to depressive symptomatology, anxiety symptomatology and suicidal ideation. *Issues Compr Pediatr Nurs*. 2007;30:87-107.
- (559) Sielk M, Altiner A, Janssen B, Becker N, de Pilars MP, Abholz HH. Prevalence and Diagnosis of Depression in Primary Care A Critical Comparison between PHQ-9 and GPs' Judgement. *Psychiatrische Praxis*. 2009;36:169-174.
- (560) Simon GE, Ludman EJ, Tutty S, Operskalski B, Von Korff M. Telephone psychotherapy and telephone care management for primary care patients starting antidepressant treatment - A randomized controlled trial. *Jama-Journal of the American Medical Association*. 2004;292:935-942.
- (561) Simon GE, Katon WJ, Lin EHB et al. Diabetes complications and depression as predictors of health service costs. *General Hospital Psychiatry*. 2005;27:344-351.
- (562) Simon GE, Ludman EJ, Linde JA et al. Association between obesity and depression in middle-aged women. *General Hospital Psychiatry*. 2008;30:32-39.
- (563) Sims J, Galea M, Taylor N et al. Regenerate: assessing the feasibility of a strength-training program to enhance the physical and mental health of chronic post stroke patients with depression. *International Journal of Geriatric Psychiatry*. 2009;24:76-83.
- (564) Sirey JA, Bruce ML, Carpenter M et al. Depressive symptoms and suicidal ideation among older adults receiving home delivered meals. *International Journal of Geriatric Psychiatry*. 2008;23:1306-1311.
- (565) Sit DKY, Flint C, Svidergol D et al. An Emerging Best Practice Model for Perinatal Depression Care. *Psychiatric Services*. 2009;60:1429-1431.
- (566) Sleath B, Chewing B, De Vellis BM et al. Communication about depression during rheumatoid arthritis patient visits. *Arthritis & Rheumatism-Arthritis Care & Research*. 2008;59:186-191.
- (567) Smith AB, Rush R, Wright P, Stark D, Velikova G, Sharpe M. Validation of an item bank for detecting and assessing psychological distress in cancer patients. *Psycho-Oncology*. 2009;18:195-199.
- (568) Smith B, Smith TC, Gray GC, Ryan MAK. When epidemiology meets the internet: Web-based surveys in the millennium cohort study. *American Journal of Epidemiology*. 2007;166:1345-1354.
- (569) Smith GC, Trauer T, Kerr PG, Chadban SJ. Prospective psychosocial monitoring of living kidney donors using the SF-36 health survey. *Transplantation*. 2003;76:807-809.
- (570) Smith MV, Poschman K, Cavaleri MA, Howell HB, Yonkers KA. Symptoms of posttraumatic stress disorder in a community sample of low-income pregnant women. *American Journal of Psychiatry*. 2006;163:881-884.

- (571) Smith TC, Smith B, Corbeil TE, Riddle JR, Ryan MAK. Self-reported mental health among US military personnel prior and subsequent to the terrorist attacks of September 11, 2001. *Journal of Occupational and Environmental Medicine*. 2004;46:775-782.
- (572) Smith TC, Smith B, Jacobson IG, Corbeil TE, Ryan MAK. Reliability of standard health assessment instruments in a large, population-based cohort study. *Annals of Epidemiology*. 2007;17:525-532.
- (573) Smith TC. The US Department of Defense Millennium Cohort Study: Career Span and Beyond Longitudinal Follow-Up. *Journal of Occupational and Environmental Medicine*. 2009;51:1193-1201.
- (574) Smolderen KG, Spertus JA, Reid KJ et al. The Association of Cognitive and Somatic Depressive Symptoms With Depression Recognition and Outcomes After Myocardial Infarction. *Circulation-Cardiovascular Quality and Outcomes*. 2009;2:328-U109.
- (575) Son YJ, Choi KS, Park YR, Bae JS, Lee JB. Depression, Symptoms and the Quality of Life in Patients on Hemodialysis for End-Stage Renal Disease. *American Journal of Nephrology*. 2009;29:36-42.
- (576) Spiegel W, Tonies H, Scherer M, Katschnig H. Learning by doing: a novel approach to improving general practitioners' diagnostic skills for common mental disorders. *Wiener Klinische Wochenschrift*. 2007;119:117-123.
- (577) Spitzer RL, Williams JBW, Kroenke K et al. Utility of new procedure for diagnosis mental-disorders in primary-care: The PRIME-MD 1000 Study. *JAMA*. 1994;272:1749-1756.
- (578) Spitzer RL, Kroenke K, Linzer M et al. Health-related quality-of-life in primary-care patients with mental disorders - Results from the PRIME-MD 1000 Study. *JAMA*. 1995;274:1511-1517.
- (579) Spitzer RL, Williams JBW, Cohen-Cole SA, Kroenke K. A new procedure for diagnosing mental disorders in primary care: The PRIME-MD Study. *Primary Psychiatry*. 1995;2:19-20.
- (580) Spitzer RL, Williams JB, Kroenke K et al. Treatment of chronic depression. *JAMA*. 1996;275:1884.
- (581) Spitzer RL, Hahn SR. PRIME-MD Revisited: A primary care option for screening mental illness. *Primary Psychiatry*. 1997;22-23.
- (582) Spitzer RL, Kroenke K, Williams JBW. Validation and utility of a self-report version of PRIME-MD - The PHQ primary care study. *JAMA*. 1999;282:1737-1744.
- (583) Spitzer RL, Williams JBW, Kroenke K, Hornyak R, McMurray J. Validity and utility of the PRIME-MD Patient Health Questionnaire in assessment of 3000 obstetric-gynecologic patients: The PRIME-MD Patient Health Questionnaire Obstetrics Gynecology Study. *Amer J Obstet Gynecol*. 2000;183:759-769.
- (584) Spitzer RL, Kroenke K, Williams JBW, Lowe B. A brief measure for assessing generalized anxiety disorder - The GAD-7. *Archives of Internal Medicine*. 2006;166:1092-1097.

- (585) Stafford L, Berk M, Jackson HJ. Validity of the Hospital Anxiety and Depression Scale and Patient Health Questionnaire-9 to screen for depression in patients with coronary artery disease. *General Hospital Psychiatry*. 2007;29:417-424.
- (586) Stauder A, Thege BK, Kovacs ME, Balog P, Williams VP, Williams RB. Worldwide Stress: Different Problems, Similar Solutions? Cultural Adaptation and Evaluation of a Standardized Stress Management Program in Hungary. *International Journal of Behavioral Medicine*. 2010;17:25-32.
- (587) Stirn A, Hinz A, Braehler E. Prevalence of tattooing and body piercing in Germany and perception of health, mental disorders, and sensation seeking among tattooed and body-pierced individuals. *Journal of Psychosomatic Research*. 2006;60:531-534.
- (588) Striegel-Moore RH, Franko DL. Epidemiology of binge eating disorder. *International Journal of Eating Disorders*. 2003;34:S19-S29.
- (589) Strine TW, Mokdad AH, Balluz LS et al. Depression and Anxiety in the United States: Findings From the 2006 Behavioral Risk Factor Surveillance System. *Psychiatric Services*. 2008;59:1383-1390.
- (590) Strine TW, Mokdad AH, Dube SR et al. The association of depression and anxiety with obesity and unhealthy behaviors among community-dwelling US adults. *General Hospital Psychiatry*. 2008;30:127-137.
- (591) Strine TW, Mokdad AH, Balluz LS, Berry JT, Gonzalez O. Impact of depression and anxiety on quality of life, health behaviors, and asthma control among adults in the United States with asthma, 2006. *Journal of Asthma*. 2008;45:123-133.
- (592) Strine TW, Dhingra SS, Kroenke K et al. Metropolitan and micropolitan statistical area estimates of depression and anxiety using the Patient Health Questionnaire-8 in the 2006 Behavioral Risk Factor Surveillance System. *International Journal of Public Health*. 2009;54:117-124.
- (593) Strine TW, Kroenke K, Dhingra S et al. The Associations Between Depression, Health-Related Quality of Life, Social Support, Life Satisfaction, and Disability in Community-Dwelling US Adults. *Journal of Nervous and Mental Disease*. 2009;197:61-64.
- (594) Sueoka K, Goulet JL, Fiellin DA et al. Depression Symptoms and Treatment Among HIV Infected and Uninfected Veterans. *Aids and Behavior*. 2010;14:272-279.
- (595) Sullivan MD, Anderson RT, Aron D et al. Health-related quality of life and cost-effectiveness components of the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial: Rationale and design. *American Journal of Cardiology*. 2007;99:901-1021.
- (596) Sundstrom IME, Bixo M, Bjorn I, Astrom M. Prevalence of psychiatric disorders in gynecologic outpatients. *Amer J Obstet Gynecol*. 2001;184:8-13.
- (597) Swindle RW, Rao JK, Helmy A et al. Integrating clinical nurse specialists into the treatment of primary care patients with depression. *International Journal of Psychiatry in Medicine*. 2003;33:17-37.

- (598) Taskapan H, Ates F, Kaya B et al. Psychiatric disorders and large interdialytic weight gain in patients on chronic haemodialysis. *Nephrology*. 2005;10:15-20.
- (599) Taylor JK, Schoenbaurn M, Katon WJ, Pincus HA, Hogan DM, Unutzer J. Strategies for identifying and channeling patients for depression care management. *American Journal of Managed Care*. 2008;14:497-504.
- (600) Thibault JM, Steiner RWP. Efficient identification of adults with depression and dementia. *American Family Physician*. 2004;70:1101-1110.
- (601) Thomas JR, Sclafani AP, Hamilton M, McDonough E. Preoperative identification of psychiatric illness in aesthetic facial surgery patients. *Aesthet Plast Surg*. 2001;25:64-67.
- (602) Thombs BD, Magyar-Russell G, Bass EB et al. Performance characteristics of depression screening instruments in survivors of acute myocardial infarction: Review of the evidence. *Psychosomatics*. 2007;48:185-194.
- (603) Thombs BD, Ziegelstein RC, Whooley MA. Optimizing Detection of Major Depression Among Patients with Coronary Artery Disease Using the Patient Health Questionnaire: Data from the Heart and Soul Study. *Journal of General Internal Medicine*. 2008;23:2014-2017.
- (604) Thombs BD, de JP, Coyne JC et al. Depression screening and patient outcomes in cardiovascular care: a systematic review. *JAMA*. 2008;300:2161-2171.
- (605) Thombs BD, Ziegelstein RC, Whooley MA. Optimizing detection of major depression among patients with coronary artery disease using the patient health questionnaire: data from the heart and soul study. *J Gen Intern Med*. 2008;23:2014-2017.
- (606) Tietjen GE, Peterlin BL, Brandes JL et al. Depression and anxiety: Effect on the migraine-obesity relationship. *Headache*. 2007;47:866-875.
- (607) Tietjen GE, Brandes JL, Digre KB et al. High prevalence of somatic symptoms and depression in women with disabling chronic headache. *Neurology*. 2007;68:134-140.
- (608) Tischler L, Schmidt S, Petermann F, Koglin U. ADHD in Adolescence - Change of Symptoms and Consequences for Research and Clinical Practice. *Zeitschrift fur Psychiatrie Psychologie und Psychotherapie*. 2010;58:23-33.
- (609) Titov N, Andrews G, Robinson E et al. Clinician-assisted Internet-based treatment is effective for generalized anxiety disorder: randomized controlled trial. *Australian and New Zealand Journal of Psychiatry*. 2009;43:905-912.
- (610) Tommasello AC, Gillis LM, Lawler JT, Bujak GJ. Characteristics of homeless HIV-positive outreach responders in urban US and their success in primary care treatment. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2006;18:911-917.
- (611) Tonn P, Reuter S, Gerlach N, Dahmen N, Pajonk FGB. Patients with psychiatric disturbances in the emergency department. *Notfall & Rettungsmedizin*. 2008;11:537-546.

- (612) Torzsa P, Rihmer Z, Gonda X et al. Family history of suicide: A clinical marker for major depression in primary care practice? *Journal of Affective Disorders*. 2009;117:202-204.
- (613) Tripp DA, Nickel JC, Ross S, Mullins C, Stechyson N. Prevalence, symptom impact and predictors of chronic prostatitis-like symptoms in Canadian males aged 16-19 years. *Bju International*. 2009;103:1080-1084.
- (614) Trivedi MH. Tools and Strategies for Ongoing Assessment of Depression: A Measurement-Based Approach to Remission. *Journal of Clinical Psychiatry*. 2009;70:26-31.
- (615) Turner AP, Williams RM, Bowen JD, Kivlahan DR, Haselkorn JK. Suicidal ideation in multiple sclerosis. *Archives of Physical Medicine and Rehabilitation*. 2006;87:1073-1078.
- (616) Turner JA, Dworkin SF. Screening for psychosocial risk factors in patients with chronic orofacial pain - Recent advances. *Journal of the American Dental Association*. 2004;135:1119-1125.
- (617) Turvey CL, Willyard D, Hickman DH, Klein DM, Kukoyi O. Telehealth screen for depression in a chronic illness care management program. *Telemedicine Journal and E-Health*. 2007;13:51-56.
- (618) Urasaki M, Oshima N, Okabayashi A et al. The Patient Health Questionnaire (PHQ-9) Scores and the Lifestyles of Nursing Students. *Southern Medical Journal*. 2009;102:800-804.
- (619) Valenstein M, Dalack G, Blow F, Figueroa S, Standiford C, Douglass A. Screening for psychiatric illness with a combined screening and diagnostic instrument. *J Gen Intern Med*. 1997;12:679-685.
- (620) Valenstein M, Kales H, Mellow A et al. Psychiatric diagnosis and intervention in older and younger patients in a primary care clinic: Effect of a screening and diagnostic instrument. *Journal of the American Geriatrics Society*. 1998;46:1499-1505.
- (621) van der Feltz-Cornelis C, Meeuwissen JAC, de Jong FJ, Hoedeman R, Elfeddali I. Randomised controlled trial of a psychiatric consultation model for treatment of common mental disorder in the occupational health setting. *Bmc Health Services Research*. 2007;7.
- (622) van der Kooy KG, van Hout HPJ, van Marwijk HWJ, de Haan M, Stehouwer CDA, Beekman ATF. Differences in heart rate variability between depressed and non-depressed elderly. *International Journal of Geriatric Psychiatry*. 2006;21:147-150.
- (623) van Marwijk HWJ, Ader H, de Haan M, Beekman A. Primary care management of major depression in patients aged ≥ 55 years: outcome of a randomised clinical trial. *British Journal of General Practice*. 2008;58:680-686.
- (624) van Ravesteijn H, Wittkamp K, Lucassen P et al. Detecting Somatoform Disorders in Primary Care With the PHQ-15. *Annals of Family Medicine*. 2009;7:232-238.
- (625) van Schaik A, van Marwijk H, Ader H et al. Interpersonal psychotherapy for elderly patients in primary care. *American Journal of Geriatric Psychiatry*. 2006;14:777-786.
- (626) VanHook MP, Berkman B, Dunkle R. Assessment tools for general health care settings: PRIME-MD, OARS, and SF-36. *Health & Social Work*. 1996;21:230-234.

- (627) Vannoy SD, Arean P, Unutzer J. Advantages of Using Estimated Depression-Free Days for Evaluating Treatment Efficacy. *Psychiatric Services*. 2010;61:160-163.
- (628) Vega WA, Rodriguez MA, Ang A. Addressing stigma of depression in Latino primary care patients. *General Hospital Psychiatry*. 2010;32:182-191.
- (629) Vogel B, Jahed J, Bengel J, Barth J, Hartner M, Baumeister H. Implementation of a Routine Gradual Psycho-Diagnostic Programme in Somatic Rehabilitation Centres - Results of a Pilot Study. *Rehabilitation*. 2009;48:361-368.
- (630) Volgsten H, Svanberg AS, Ekselius L, Lundkvist O, Poromaa IS. Prevalence of psychiatric disorders in infertile women and men undergoing in vitro fertilization treatment. *Human Reproduction*. 2008;23:2056-2063.
- (631) Vuorilehto M, Melartin T, Isometsa E. Depressive disorders in primary care: recurrent, chronic, and co-morbid. *Psychological Medicine*. 2005;35:673-682.
- (632) Vuorilehto MS, Melartin TK, Isometsa ET. Suicidal behaviour among primary-care patients with depressive disorders. *Psychological Medicine*. 2006;36:203-210.
- (633) Vuorilehto MS, Melartin TK, Isometsa ET. Course and outcome of depressive disorders in primary care: a prospective 18-month study. *Psychological Medicine*. 2009;39:1697-1707.
- (634) Walker J, Waters RA, Murray G et al. Better off dead: Suicidal thoughts in cancer patients. *Journal of Clinical Oncology*. 2008;26:4725-4730.
- (635) Walker J, Hansen CH, Hodges L et al. Screening for suicidality in cancer patients using Item 9 of the nine-item patient health questionnaire; does the item score predict who requires further assessment? *General Hospital Psychiatry*. 2010;32:218-220.
- (636) Watson LC, Zimmerman S, Cohen LW, Dominik R. Practical Depression Screening in Residential Care/Assisted Living: Five Methods Compared With Gold Standard Diagnoses. *American Journal of Geriatric Psychiatry*. 2009;17:556-564.
- (637) Weber T, Keller M, Hense I et al. Effect of parathyroidectomy on quality of life and neuropsychological symptoms in primary hyperparathyroidism. *World Journal of Surgery*. 2007;31:1202-1209.
- (638) Weiss BD, Francis L, Senf JH, Heist K, Hargraves R. Literacy education as treatment for depression in patients with limited literacy and depression: A randomized controlled trial. *Journal of General Internal Medicine*. 2006;21:823-828.
- (639) Weissman MM, Feder A, Pilowsky DJ et al. Depressed mothers coming to primary care: maternal reports of problems with their children. *Journal of Affective Disorders*. 2004;78:93-100.
- (640) Weobong B, Akpalu B, Doku V et al. The comparative validity of screening scales for postnatal common mental disorder in Kintampo, Ghana. *Journal of Affective Disorders*. 2009;113:109-117.

- (641) Wessels AM, Pouver F, Geelhoed-Duijvestijn PHLM et al. No evidence for increased self-reported cognitive failure in Type 1 and Type 2 diabetes: a cross-sectional study. *Diabetic Medicine*. 2007;24:735-740.
- (642) West CP, Tan AD, Habermann TM, Sloan JA, Shanafelt TD. Association of Resident Fatigue and Distress With Perceived Medical Errors. *Jama-Journal of the American Medical Association*. 2009;302:1294-1300.
- (643) Whooley MA, Avins AL, Miranda J, Browner WS. Case-finding instruments for depression. Two questions are as good as many. *J Gen Intern Med*. 1997;12:439-445.
- (644) Whooley MA. Depression and cardiovascular disease: healing the broken-hearted. *JAMA*. 2006;295:2874-2881.
- (645) Whooley MA, Caska CM, Hendrickson BE, Rourke MA, Ho J, Ali S. Depression and inflammation in patients with coronary heart disease: findings from the Heart and Soul Study. *Biol Psychiatry*. 2007;62:314-320.
- (646) Whooley MA, De Jonge P, Vittinghoff E et al. Depressive Symptoms, Health Behaviors, and Risk of Cardiovascular Events in Patients With Coronary Heart Disease. *Jama-Journal of the American Medical Association*. 2008;300:2379-2388.
- (647) Whooley MA, de JP, Vittinghoff E et al. Depressive symptoms, health behaviors, and risk of cardiovascular events in patients with coronary heart disease. *JAMA*. 2008;300:2379-2388.
- (648) Williams JBW, Spitzer RL, Linzer M et al. Gender differences in depression in primary-care. *Amer J Obstet Gynecol*. 1995;173:654-659.
- (649) Williams LS, Plue L, Brizendine E, Tu WZ, Hendrie H, Kroenke K. Performance of the PHQ-9 as a screening tool for post-stroke depression. *Stroke*. 2004;35:321.
- (650) Williams LS, Jones WJ, Shen J, Robinson RL, Kroenke K. Outcomes of newly referred neurology outpatients with depression and pain. *Neurology*. 2004;63:674-677.
- (651) Williams LS, Brizendine EJ, Plue L et al. Performance of the PHQ-9 as a screening tool for depression after stroke. *Stroke*. 2005;36:635-638.
- (652) Williams LS, Kroenke K, Bakas T et al. Care management of poststroke depression - A randomized, controlled trial. *Stroke*. 2007;38:998-1003.
- (653) Williams RT, Heinemann AW, Bode RK, Wilson CS, Fann JR, Tate DG. Improving Measurement Properties of the Patient Health Questionnaire-9 With Rating Scale Analysis. *Rehabilitation Psychology*. 2009;54:198-203.
- (654) Wilson KG, Chochinov HM, Skirko MG et al. Depression and anxiety disorders in palliative cancer care. *Journal of Pain and Symptom Management*. 2007;33:118-129.
- (655) Wilson MW, Richards JS, Klapow JC, Devivo MJ, Greene P. Cluster analysis and chronic pain: An empirical classification of pain subgroups in a spinal cord injury sample. *Rehabilitation Psychology*. 2005;50:381-388.

- (656) Wittkamp K, van Ravesteijn H, Baas K et al. The accuracy of Patient Health Questionnaire-9 in detecting depression and measuring depression severity in high-risk groups in primary care. *General Hospital Psychiatry*. 2009;31:451-459.
- (657) Wittkamp KA, Naeije L, Schene AH, Huyser J, van Weert HC. Diagnostic accuracy of the mood module of the Patient Health Questionnaire: a systematic review. *General Hospital Psychiatry*. 2007;29:388-395.
- (658) Woods SM, Melville JL, Guo YQ, Fan MY, Gavin A. Psychosocial stress during pregnancy. *American Journal of Obstetrics and Gynecology*. 2010;202.
- (659) Wryobeck JM, Walton MA, Curran GM, Massey LS, Booth BM. Complexities of cocaine users presenting to the emergency department with chest pain: Interactions between depression symptoms, alcohol use, and race. *Journal of Addiction Medicine*. 2007;1:213-221.
- (660) Wulsin L, Somoza E, Heck J. The Feasibility of Using the Spanish PHQ-9 to Screen for Depression in Primary Care in Honduras. *Prim Care Companion J Clin Psychiatry*. 2002;4:191-195.
- (661) Wulsin LR, Musselman D, Otte C, Bruce E, Ali S, Whooley MA. Depression and whole blood serotonin in patients with coronary heart disease from the Heart and Soul Study. *Psychosom Med*. 2009;71:260-265.
- (662) Yawn BP, Pace W, Wollan PC et al. Concordance of Edinburgh Postnatal Depression Scale (EPDS) and Patient Health Questionnaire (PHQ-9) to Assess Increased Risk of Depression among Postpartum Women. *Journal of the American Board of Family Medicine*. 2009;22:483-491.
- (663) Yeung A, Yu SC, Fung F, Vorono S, Fava M. Recognizing and engaging depressed Chinese Americans in treatment in a primary care setting. *International Journal of Geriatric Psychiatry*. 2006;21:819-823.
- (664) Yeung A, Fung F, Yu SC et al. Validation of the Patient Health Questionnaire-9 for depression screening among chinese Americans. *Comprehensive Psychiatry*. 2008;49:211-217.
- (665) Yonkers KA, Smith MV, Lin HQ, Howell HB, Shao L, Rosenheck RA. Depression Screening of Perinatal Women: An Evaluation of the Healthy Start Depression Initiative. *Psychiatric Services*. 2009;60:322-328.
- (666) Zaharna AF, Budur K, Hagan J, Gonsalves L. Association between sleep disturbances and suicidal thoughts in women with depression: A cross-sectional survey using PHQ-9. *Sleep*. 2008;31:959.
- (667) Zaider TI, JOHNSON JG, Cockell SJ. Psychiatric comorbidity associated with eating disorder symptomatology among adolescents in the community. *Int J Eat Disord*. 2000;28:58-67.
- (668) Zaider TI, JOHNSON JG, Cockell SJ. A prospective longitudinal investigation of psychiatric disorders associated with onset and persistence of eating disorders during adolescence. *Journal of Youth and Adolescence*. 2002;33:319-329.

- (669) Zane RD, McAfee AT, Sherburne S, Billeter G, Barsky A. Panic disorder and emergency services utilization. *Academic Emergency Medicine*. 2003;10:1065-1069.
- (670) Zastrow A, Faude V, Seyboth F, Niehoff D, Herzog W, Lowe B. Risk factors of symptom underestimation by physicians. *Journal of Psychosomatic Research*. 2008;64:543-551.
- (671) Zuckerbrot RA, Jensen PS. Improving recognition of adolescent depression in primary care. *Arch Pediatr Adolesc Med*. 2006;160:694-704.